ABNORMAL INVOLUNTARY MOVEMENT SCALE (AIMS)

Public Health Service	NAME:
Alcohol, Drug Abuse, and Mental Health Administration	DATE:
National Institute of Mental Health	Prescribing Practitioner:

CODE: 0 = None

1 = Minimal, may be extreme normal

INSTRUCTIONS:2 = MildComplete Examination Procedure (attachment d.)3 = Moderatebefore making ratings4 - Severe

MOVEMENT RATINGS: Rate highest severity observed. Rate		RATER	RATER		RATER		RATER		RATER	
		upon activation one <u>less</u> than those observed								
spontaneously. Circle movement as well as code number that		Date		Date		Date		Date		
applies.										
Facial and	1.	Muscles of Facial Expression	0 1 2 3	3 4	0 1 2	2 3 4	0 1 2	2 3 4	0 1 2	2 3 4
Oral		e.g. movements of forehead, eyebrows								
Movements		periorbital area, cheeks, including frowning								
		blinking, smiling, grimacing								
	2.	Lips and Perioral Area	0 1 2 3	3 4	0 1 2	2 3 4	0 1 2	2 3 4	0 1 2	2 3 4
		e.g., puckering, pouting, smacking								
	3.	Jaw e.g. biting, clenching, chewing, mouth	0 1 2 3	3 4	0 1 2	2 3 4	0 1 2	2 3 4	0 1 2	2 3 4
		opening, lateral movement								
	4.	Tongue Rate only increases in movement								
		both in and out of mouth. NOT inability to	0 1 2 3	3 4	0 1 2	2 3 4	0 1 2	2 3 4	O 1	2 3 4
		sustain movement. Darting in and out of								
		mouth.								
	5.	Upper (arms, wrists,, hands, fingers)								
		Include choreic movements (i.e., rapid,								
Extremity		objectively purposeless, irregular,								
Movements		spontaneous) athetoid movements (i.e., slow,	0 1 2 3	3 4	0 1 2	2 3 4	0 1 2	2 3 4	0 1 2	2 3 4
		irregular, complex, serpentine). DO NOT								
		INCLUDE TREMOR (i.e., repetitive,								
		regular, rhythmic)								
	6.	Lower (legs, knees, ankles, toes)								
		e.g., lateral knee movement, foot tapping,								
		heel dropping, foot squirming, inversion and	0 1 2 3	3 4	0 1 2	2 3 4	0 1 2	2 3 4	0 1 2	2 3 4
		eversion of foot.								
Trunk	7.	Neck, shoulders, hips e.g., rocking,	0 1 2	3 4	0 1 2	2 3 4	0 1 2	2 3 4	0 1 2	2 3 4
Movements		twisting, squirming, pelvic gyrations								
	8.	Severity of abnormal movements overall	0 1 2 3	3 4	0 1 2	2 3 4	0 1 2	2 3 4	0 1 2	2 3 4
Global	9.	Incapacitation due to abnormal	0 1 2	3 4	0 1 2	2 3 4	0 1 2	2 3 4	0 1 2	2 3 4
Judgments		movements								
8	10.	Patient's awareness of abnormal								
		movements. Rate only patient's report								
		No awareness 0	0 1 2 3		0 1 2 3		0 1 2 3		0	
		Aware, no distress 1							1	
		Aware, mild distress 2								2
		Aware, moderate distress 3								3
		Aware, severe distress 4		4		4		4		4
	11.	Current problems with teeth and/or								
Dental Status		dentures	No	Yes	No	Yes	No	Yes	No	Yes
				Yes	No	Yes	No	Yes	No	Yes
	12.	Are dentures usually worn?		-						
		, , , , , , , , , , , , , , , , , ,	No	Yes	No	Yes	No	Yes	No	Yes
	13.	Edentia?								
	1		No	Yes	No	Yes	No	Yes	No	Yes
	14.	Do movements disappear in sleep?		1 30	1.0	- 00	1,0	- 05	1,0	. 00
	1 1.	20 moremento albappeni in bicepi	1		L		l		<u> </u>	

Final: 9/2000