telpful HEDIS Documentation Tips for Pediatric Providers

HEDIS Measure Definitions	What You Can Do	Coding/Tips
Well Child Visits — 3 age groups:	Never miss an opportunity! Exam require- ments can be performed during a sick visit or a well-child exam.	ICD-10 CM Codes: Z00.11 -Z00.129, Z00.5, Z00.8, Z02.0-Z02.9, Z76.1- Z76.2, Z00.00-Z00.01
 W15 - Well Child 15 months Members 0-15 months of age with 6 comprehensive well child visits. Minimum of 6 well visits required by 15 months old W34 - Well Child 3-6 years Members 3-6 years of age with at least 1 comprehensive well child visits annually. Minimum of 1 visit required annually AWC - Adolescent Well Care Visits Members 12-21 years of age with at least one comprehensive well care visit with a primary care practitioner or an OB/GYN practitioner annually. Minimum of 1 required annually 	 Documentation MUST include ALL of the following: A health history – assessment of member's history of disease or illness and family health history A physical development history- assessment of specific age appropriate physical development milestones A mental development history – assessment of specific age-appropriate mental development milestones A physical exam Health education/anticipatory guidance – guidance given in anticipation of emerging issues that a child/family may face 	 CPT Codes: 99381—99385, 99391 - 99395, 99461 HCPCS: G0438, G0439 Documentation that Does NOT count as compliant: For Health History: notation of allergies or medications or immunization status alone. If all three are documented it meets health history For Physical Development History: notation of appropriate for age without specific mention of development; notation of well-developed/ nourished; tanner stage (except for adolescents—then it meets compliance) For Mental Development History: notation of appropriately responsive for age; neurological exam; notation of well-developed For Physical Exam—vital signs alone; for adolescent visits to an OB/GYN they do not meet compliance if the visit is limited to OB/GYN topics For Health Education/Anticipatory Guidance - information regarding medications or immunizations or their side effects. Handouts given
WCC - Weight Assessment and Counseling for Nutrition and Physical Activity for Chil- dren/Adolescents Children age 3-17 years of age who had a visit with a PCP or OB/GYN and who had BMI percentile documentation, and counsel- ing for nutrition and physical activity	Document height, weight and BMI <u>percentile.</u> Discussion and documentation of nutrition and physical activity during at least one office visit annually. Examples Nutrition— discussion of current nutrition behaviors; weight or obesity counseling Physical Activity—discussion of current physi- cal activity behaviors, exercise routine, sports activities; sports physical, weight or obesity counseling	during a visit without evidence of discussion.BMI_ICD-10 CM Codes:Z68.51-Z68.54Nutrition Counseling -ICD-10 CM Code:Z71.3CPT Codes:97802-97804HCPCS:G0447,G0270,G0271,S9449,S9452,S9470Physical Activity CounselingICD-10 CM Code:Z02.5 (Sports physical)Z71.82(Exercise counseling)
 IMA - Immunizations in Adolescents Members who turned 13 years of age in the measurement year and received by age 13: Tdap vaccine—one dose between the 10th and 13th birthday Meningococcal Conjugate vaccine - one dose of meningococcal serogroups A,C,W, Y vaccine between the 11th and 13th birthday HPV—either two doses of HPV vaccine between the 9th and 13th birthday with at least 146 days between doses OR three dose os with different dates of service between the 9th and 13th birthday. 	Educate staff to schedule PRIOR to 13th birth- day. Give call reminders for series vaccines Meningococcal recombinant (serogroup B) vaccines <u>Do Not Count</u> . Be sure your immun- ization claims and records are clear about which meningococcal was given! Document and submit claims timely with correct code. HPV rates are now reported for both females and males. Educate families on the importance of these immunizations.	Tdap CPT Code: 90715 CVX Code: 115 Meningococcal CPT Codes: 90734 CVX Codes: 108, 114, 136, 147, 167 HPV CPT Codes: 90649, 90650, 90651 CVX Codes: 62, 118, 137, 165

HEDIS Measure Definitions	What You Can Do	Coding/Tips
CIS/LCS - Childhood Immunization Status and Lead Screening in Children Children who received recommended vaccinations prior to second birthday.	Educate office staff to schedule appoint- ments PRIOR to 2nd birthday. Call families to schedule appointments for those that are behind with their immunizations. Any vaccines after the age of 2 are consid-	Vaccine Codes DTaP CPT Codes: 90698, 90700, 90721, 90723 CVX Codes: 20, 50, 106, 107, 110, 120 IPV CPT Codes: 90698, 90713, 90723
Children who had one or more lead blood test for lead poisoning by their second birthday. *Document parental refusal. *	 ered late in HEDIS reporting. Educate parents/guardians regarding the importance of having their child immunized and keeping appointments. Immunizations recommended: 4 DTaP/DT, 3 IPV, 1 MMR, 3 HiB, 3 Hep B, 1 VZV, 4 PCV, 1 Hep A, 2 or 3 Rotavirus and 2 Influenza vaccines by the second birthday. Document in the medical record if member has evidence of the disease for which immunization is intended or if a contraindication to the vaccine exists. Lead screening test should be completed on all children before their second birthday. A lead risk questionnaire does not count - it must be a capillary or venous blood lead test. 	CVX Codes: 10, 89, 110, 120 HiB CPT Codes: 90644-90648, 90698, 90721, 90748 CVX Codes: 17, 46 –51, 120, 148 HepB CPT Codes: 90723, 90740, 90744, 90747, 90748 CVX Codes: 08, 44, 45, 51, 110 HCPCS:G0010 PCV CPT Codes: 90670 CVX Codes: 133, 152 HCPCS: G0009 VZV CPT Codes: 90710, 90716 CVX Codes: 21, 94 MMR CPT Codes: 90707, 90710 CVX Codes: 03, 94 Measles CPT Code: 90705 CVX Code: 05 Measles/Rubella CPT Code: 90708 CVX Code: 04 Mumps CPT Code: 90704 CVX Code: 07 Rubella CPT Code: 90706 CVX Code: 119 Rotavirus 2 dose CPT Code: 90681 CVX Code: 119 Rotavirus 3 dose CPT Code: 90680 CVX Code: 116,122 HepA CPT Code: 90655, 90657, 90661, 90662, 90673, 90685 -90689 HCPCS: G0008 CVX Codes: 88, 135, 140, 141, 150, 153, 155, 158, 161
		Live Attenuated influenza (Nasal): CVX Code: 111, 149 CPT Code: 90660, 90672 Lead Screening CPT Code: 83655
 ADD - Follow-Up Care for Children Prescribed ADHD Medication Children 6-12 years of age, newly prescribed ADHD medication who had at least 3 follow-up visits within a 10 month period, one of which was within 30 days of when the ADHD medication was dispensed. Two rates are reported: Initiation Phase: A follow-up visit with a practitioner with prescribing authority during the 30 day initiation phase Continuation Phase: children that remained on the ADHD medication for at least 210 days, and in addition to the visit in the Initiation Phase, had at least 2 follow -up visits with a practitioner within 270 days (9 months) after the Initiation Phase ended. 	 When prescribing a new ADHD medication for a patient, schedule the initial follow-up appointment before the patient leaves the office. Schedule the initial follow-up for 2-3 weeks after starting the medication. Explain to the parent/guardian the im- portance of follow-up care No refills unless the child has the initial follow-up visit After the initial follow-up visit, schedule at least 2 more visits over the next 9 months to check the child's progress Encourage parents/caregivers to ask ques- tions about their child's ADHD 	BH Stand Alone OP Visit Codes CPT : 98960-98962, 99078, 99201-99205, 99211-99215, 99241-99245, 99341-99345, 99347-99350, 99381-99387, 99391-99397, 99401-99404, 99411, 99412, 99483, 99510 HCPCS: G0155, G0176, G0177, G0409, G0463, H0002, H0004, H0031, H0034, H0036-H0037, H0039, H0040, H2000, H2010-H2011, H2013-H220, M0064, T1015 UB REV: 0510, 0513, 0515-0517, 0519-0523, 0526-0529, 0900, 0902-0904, 0911, 0914 -0917, 0919, 0982, 0983 Observation Visit CPT Codes: 99217-99220 Health & Behavior Assessment/Intervention CPT Codes: 96150-96154 Intensive OP encounter/Partial Hospitalization Codes HCPCS: G0410-0411, H0035, H2001, H2012, S0201, S9480, S9484-9485 UBREV: 905, 907, 912, 913 CPT codes that require a POS code:
	One follow-up visit in the Continuation phase can also be completed via telephone Telephone Visit CPT Codes: 98966-98968, 99441-99443 or a Telehealth Modifier: 95, GT or Telehealth POS: 2 added to a visit code	CPT : 90791-90792, 90832-90834, 90836-90840, 90845, 90847, 90849, 90853, 90875, 90876, 99221-99223, 99231- 99233, 99238, 99239, 99251-99255 POS : 02. 03, 05, 07, 09, 11-20, 22, 33, 49, 50, 52, 53, 71, 72 POS 2 (telehealth) permitted for only one of the follow -up visits in the continuation phase)

HEDIS Measure Definitions	What You Can Do	Coding/Tips
URI - Appropriate Treatment for Upper Respiratory Infection	Do not prescribe antibiotics for URI treat- ment.	ICD-10 CM Codes : J00, J06.0, J06.9
Members age 3 months and older with a diagnosis of upper respiratory infection (URI) and that did NOT result in an antibi- otic dispensing event.	Document and submit all appropriate diagnosis on claims if more than one diagnosis is appropriate. A competing diagnosis of pharyngitis or other infec- tion on the same date or 3 days after will exclude the member.	
CWP - Appropriate Testing for Children with Pharyngitis	Before prescribing an antibiotic for a diagnosis of pharyngitis , perform a group A strep test.	Pharyngitis ICD-10 CM Codes: J02.0, J02.8-J03.01, J03.80- J03.81, J03.90-J03.91
Members age 3 years and older where the member was diagnosed with pharyngitis, dispensed and antibiotic and received a group A strep test for the episode.	Document and submit claims for all ap- propriate diagnoses established at the visit Submit claim for in-office rapid strep test	Group A Strep Tests CPT Codes: 87070, 87071, 87081, 87430, 87650-87652, 87880
	There are numerous comorbid condition and competing diagnoses exclusions for this measure.	
AAB - Avoidance of Antibiotic Treatment for Acute Bronchitis/Bronchiolitis	Treat acute bronchitis primarily with home treatments to relieve symptoms. Antibiotics don't usually help (viral).	Acute Bronchitis or Bronchiolitis: ICD-10 CM Codes: J20.3-J20.9, J21.0-J21.1, J21.8-J21.9
Members age 3 months and older with a diagnosis of acute bronchitis/bronchiolitis that did not results in an antibiotic dispens- ing event. This measure used to be for adults only and now includes everyone ages 3 months and older	Of course, some patients have comorbid conditions and require antibiotics. These patients would be excluded from this measure reporting. A diagnosis of phar- yngitis on the same day or in the 3 days after also exclude this member. Educate patients about overuse of antibi- otics and resistance.	
CHL - Chlamydia Screening in Women	Assist with member education about STDs, transmission and the importance of testing.	CPT Codes: 87110, 87270, 87320, 87490-87492, 87810
Women 16-24 years of age who are identi- fied as sexually active and had at least one Chlamydia test annually.	Perform routine urine test for Chlamydia, document and submit claims timely.	
MMA– Medication Management for Peo- ple With Asthma	Schedule regular follow-up for people with persistent asthma	Asthma Controller Medications
Members age 5-64, identified as having persistent asthma and dispensed appropri- ate medications that they remained on during the treatment period (end of calen- dar year)	Patient education about benefits of med- ication compliance Order medications that are on the member's health plan formulary	Antiasthmatic Combinations - Dyphylline-guaifenesin Antibody Inhibitor - Omalizumab Anti-interleukin-5— Benralizumab, Mepolizumab, Reslizumab Inhaled Steroid Combinations - Budesonise-formoterol, Formoterol-mometasone, Fluticasone-salmeterol,
 Remained on asthma controller medication for at least 50% of the treatment period. 	Exclusions—anytime in patient's history Acute Respiratory Failure ICD-10: J96.00 -J96.02, J96.20-J96.22 Chronic Respiratory Conditions due to Fumes/Vapors ICD-10: J68.4 COPD ICD-10: J44.0, J44.1, J44.9	Fluticasone-vilanterol Inhaled Corticosteroids - Beclomethasone, Budesonise, Ciclesonide, Flunisolide, Fluticasone, Mometasone Leukotriene Modifiers - Montelukast, Zafirlukast, Zileuton Methylxanthines - Aminophylline, Theophylline Asthma Reliever Medications
 Remained on asthma controller medi- cation for at least 75% of the treat- ment period. 	Cystic Fibrosis ICD-10: E84.0, E84.11, E84.19, E84.8, E84.9 Emphysema ICD-10: J43.0-J43.2, J43.8- J43.9 Other Emphysema ICD-10: J98.2, J98.3	Short-acting, inhaled beta-2 agonists - Albuterol, Levalbut- erol
ABA - Adult BMI Assessment	Perform and document criteria of Ht/Wt/ BMI calculation at each visit or at least	ICD-10 CM Codes:
Documentation of body mass index (BMI) and weight annually or every other year in members 18—74 years of age.	annually. Patients younger than 20 years old need to have a BMI percentile documented *Pregnant members are excluded from this measure*	BMI - Z68.1, Z68.20-Z68.29, Z68.30-Z68.39, Z68.41-Z68.45 BMI Percentile - Z68.51-Z68.54

HEDIS Measure Definitions	What You Can Do	Coding/Tips
 CDC - Comprehensive Diabetes Care Members 18-75 years of age with diabetes should have each of the following: HbA1C testing, HbA1C control Medical attention for nephropathy Retinal eye exam (refer for exam) Blood pressure control 	Order screenings annually or more often as needed and educate members on importance of compliance with testing and medications. Include all current medications on the medication list. Be sure to indicate if a member is on an ACE/ARB medication Refer member to Optometrist or Oph- thalmologist for Dilated Retinal Eye Exam annually. Explain why this is important and that it is different than an eye for glasses or contacts. Document Stage 4 chronic kidney disease or End State Renal Disease (ESRD) with appropriate codes: Stage 4 chronic kidney disease ICD-10 CM: N18.4 ESRD ICD-10 CM: N18.5, N18.6, Z91.15, Z99.2	Diabetes ICD-10 CM Codes: E10.10-E13.9, O24.011-O24.33, O24.811-O24.83 HbA1c CPT Codes: 83036, 83037 CPTII HbA1C Result Codes HbA1c level less than 7.0: 3044F HbA1c level greater than 9.0: 3046F HbA1c level greater > or = 7 & < 8: 3051F HbA1c level greater > or = 8 & < 9: 3052F Urine Protein Tests - check annually, especially if not on an ACE/ARB medication CPT Codes: 81000-81003, 81005 , 82042-82044, 84156 CPT II Codes: 3060F-3062F Blood Pressure CPT Codes: Systolic BP: < 130 3074F, 130-139 3075F ; >/= to 140 3077F Diastolic BP: 80-89 3079F; < 80 3078F; >/= 90 3080F
CBP - Controlling High Blood Pressure	If BP elevated (140/90 or greater) at initial vital sign assessment, alleviate	ICD-10 CM Code: 110 Blood Pressure CPT Codes:
 Members 18-85 years of age with a diagnosis of hypertension (HTN) and whose BP is adequately controlled (<140/90). Blood pressure value CPT II codes are acceptable to meet compliance BP readings taken from an electronic device and results were digitally stored and transmitted to the provider and interpreted by the provider, can be used Antidepressant Medication Management (AMM) Members 18 years of age and older who were treated with antidepressant medication, had a diagnosis of major depression and who remained on antidepressant medication treatment. Two rates are reported: Effective Acute Phase: Percentage of patients who remained on an antidepressant medication for at least 84 days (12 weeks) Effective Continuation Phase: Percentage of patients who remained on an antidepressant medication for at least 180 days (6 months) 	potential factors that might cause temporary elevation and retake BP during exam. Make sure you use the correct size cuff. If using a automatic BP machine, record the actual number, do NOT round up. Schedule follow up visits to monitor effectiveness of BP medication. Educate members that medication may take several weeks to become effective, they should call with any potential medication concerns/reactions Stress that they should not stop medication abruptly or without consulting you first for assistance Schedule follow up appointments prior to patient leaving your office Outreach patients that cancel appointments and have not rescheduled Stress the importance of medication compliance.	Systolic BP: < 130 3074F, 130-139 3075F; >/= to 140 3077F Diastolic BP: 80-89 3079F; < 80 3078F; >/= 90 3080F Exclusions: End Stage Renal Disease (ESRD) or a kidney transplant on or prior to December 31st of the measurement year or a diag- nosis of pregnancy during the measurement year, would exclude someone from this measure. ICD-10 CM Codes for Major Depression: F32.0-F32.4, F32.9, F33.0-F33.3, F33.41, F33.9
ADV—Annual Dental Visit Members 2-20 years of age who had at least on dental visit during the measurement year.	Educate parents/guardians about the importance of brushing from an early age as well as dental visits as early as age 2 Ask when the last dental visit was and remind them to schedule one if they have not been.	Any claim with a dental practitioner during the measure- ment year meets compliance.