



AETNA BETTER HEALTH®  
Coverage Policy/Guideline

Name: Dupixent

Page: 1 of 16

Effective Date: 10/16/2025

Last Review Date: 9/9/2025

Applies to:  Illinois

### Intent:

The intent of this policy/guideline is to provide information to the prescribing practitioner outlining the coverage criteria for Dupixent under the patient's prescription drug benefit.

### Description:

The indications below including FDA-approved indications and compendial uses are considered a covered benefit provided that all the approval criteria are met, and the member has no exclusions to the prescribed therapy.

#### FDA-approved Indications

- Treatment of adult and pediatric patients aged 6 months and older with moderate-to-severe atopic dermatitis whose disease is not adequately controlled with topical prescription therapies or when those therapies are not advisable. Dupixent can be used with or without topical corticosteroids.
- Add-on maintenance treatment of adult and pediatric patients aged 6 years and older with moderate-to-severe asthma characterized by an eosinophilic phenotype or with oral corticosteroid dependent asthma.
- Add-on maintenance treatment in adult and pediatric patients aged 12 years and older with inadequately controlled chronic rhinosinusitis with nasal polyps (CRSwNP).
- Treatment of adult and pediatric patients aged 1 year and older, weighing at least 15 kg, with eosinophilic esophagitis (EoE).
- Treatment of adult patients with prurigo nodularis (PN).
- Add-on maintenance treatment of adult patients with inadequately controlled chronic obstructive pulmonary disease (COPD) and an eosinophilic phenotype.
- Treatment of adult and pediatric patients aged 12 years and older with chronic spontaneous urticaria (CSU) who remain symptomatic despite H1 antihistamine treatment.
- Treatment of adult patients with bullous pemphigoid (BP)

#### Limitations of Use

- Not indicated for the relief of acute bronchospasm or status asthmaticus.
- Not indicated for treatment of other forms of urticaria.

#### Compendial Uses

Immune checkpoint inhibitor-related toxicities

All other indications are considered experimental/investigational and not medically necessary.

### Applicable Drug List:

Dupixent



AETNA BETTER HEALTH®  
Coverage Policy/Guideline

Name: Dupixent

Page: 2 of 16

Effective Date: 10/16/2025

Last Review Date: 9/9/2025

Applies to:  Illinois

## Policy/Guideline:

### Documentation:

Submission of the following information is necessary to initiate the prior authorization review:

### Atopic dermatitis

#### Initial requests:

Chart notes, medical record documentation, or claims history supporting previous medications tried, including response to therapy. If therapy is not advisable, documentation of clinical reason to avoid therapy.

#### Continuation requests:

Provider attestation that the member has experienced a positive clinical response to therapy as evidenced by low disease activity or improvement in signs or symptoms of atopic dermatitis.

### Asthma

#### Initial requests:

- Chart notes or medical record documentation showing baseline blood eosinophil count (where applicable).
- Chart notes, medical record documentation, or claims history supporting previous medications tried including drug, dose, frequency, and duration.

#### Continuation requests:

Provider attestation supporting improvement in asthma control.

### Chronic rhinosinusitis with nasal polyps (CRSwNP)

#### Initial requests:

Chart notes, medical record documentation, or claims history supporting previous medications tried. If therapy is not advisable, documentation of clinical reason to avoid therapy.

#### Continuation requests:

Provider attestation supporting positive clinical response.

### Eosinophilic esophagitis (EoE)

#### Initial requests:

- Chart notes or medical record documentation showing endoscopic biopsy details including intraepithelial esophageal eosinophil count.
- Chart notes, medical record documentation, or claims history supporting previous medications tried. If therapy is not advisable, documentation of clinical reason to avoid therapy.

#### Continuation requests:

Chart notes or medical record documentation supporting positive clinical response.



AETNA BETTER HEALTH®  
Coverage Policy/Guideline

Name: Dupixent

Page: 3 of 16

Effective Date: 10/16/2025

Last Review Date: 9/9/2025

Applies to:  Illinois

## **Prurigo Nodularis (PN)**

### Initial requests

- Chart notes or medical record documentation of symptoms (e.g., pruritus, nodular lesions) (where applicable).
- Chart notes, medical record documentation, or claims history supporting previous therapies tried, including response to therapy. If therapy is not advisable, documentation of clinical reason to avoid therapy.

### Continuation requests

Chart notes or medical record documentation supporting positive clinical response.

## **Chronic Obstructive Pulmonary Disease (COPD)**

### Initial requests

- Chart notes or medical record documentation demonstrating classic signs and/or symptoms of COPD.
- Chart notes, medical record documentation, or claims history of previous medications tried, including response to therapy. If therapy is not advisable, documentation of clinical reason to avoid therapy.
- Chart notes or medical record documentation showing absolute blood eosinophil count prior to initiating therapy with the requested medication.
- Chart notes or medical record documentation of moderate or severe exacerbations within the last year.

### Continuation requests

Chart notes or medical record documentation supporting positive clinical response.

## **Chronic Spontaneous Urticaria (CSU)**

### Initial requests

Chart notes, medical record documentation, or claims history supporting previous medications tried, including response to therapy.

### Continuation requests

Chart notes or medical record documentation supporting positive response to therapy.

## **Immune Checkpoint Inhibitor-Related Toxicities**

### Continuation requests

Chart notes or medical record documentation supporting positive clinical response.

## **Bullous Pemphigoid (BP)**

### Initial requests

- Chart notes or medical record documentation demonstrating clinical features of bullous pemphigoid.



AETNA BETTER HEALTH®  
Coverage Policy/Guideline

Name: Dupixent

Page: 4 of 16

Effective Date: 10/16/2025

Last Review Date: 9/9/2025

Applies to:  Illinois

- Chart notes, medical record documentation, or claims history of previous medications tried, including response to therapy. If therapy is not advisable, documentation of clinical reason to avoid therapy.

#### Continuation requests

Chart notes or medical record documentation supporting positive clinical response.

#### **Prescriber Specialties:**

This medication must be prescribed by or in consultation with ONE of the following:

- Atopic dermatitis: any
- Asthma: allergist/immunologist or pulmonologist
- Chronic rhinosinusitis with nasal polyps: allergist/immunologist or otolaryngologist
- Eosinophilic esophagitis: gastroenterologist or allergist/immunologist
- Prurigo nodularis: dermatologist or allergist/immunologist
- Chronic obstructive pulmonary disease: pulmonologist or allergist/immunologist
- Chronic spontaneous urticaria: allergist/immunologist or dermatologist
- Immune checkpoint inhibitor-related toxicity and bullous pemphigoid: dermatologist, hematologist or oncologist

#### **Coverage Criteria:**

##### **Atopic dermatitis**

Authorization of 6 months may be granted for members 6 months of age or older who have previously received a biologic or targeted synthetic drug indicated for moderate-to-severe atopic dermatitis in the past year.

OR

Authorization of 6 months may be granted for treatment of moderate-to-severe atopic dermatitis in members 6 months of age or older when both of the following criteria are met:

- Member has had an inadequate treatment response with a medium to high potency topical corticosteroid; AND
- Member has an inadequate response with ONE of the following in the past 2 years:
  - Generic immunosuppressant
  - Topical Calcineurin Inhibitors (TCI)
  - Phototherapy
  - Phosphodiesterase-4 inhibitor (PDE-4)

##### **Asthma**

Authorization of 6 months may be granted for members 6 years of age or older who have previously received a biologic drug indicated for asthma in the past year.

OR



AETNA BETTER HEALTH®  
Coverage Policy/Guideline

Name:	Dupixent	Page:	5 of 16
Effective Date:	10/16/2025	Last Review Date:	9/9/2025
Applies to:	<input checked="" type="checkbox"/> Illinois		

Authorization of 6 months may be granted for treatment of moderate-to-severe asthma in members 6 years of age or older when ONE of the following criteria are met:

- Member has a baseline blood eosinophil count of at least 150 cells per microliter and 1 exacerbation (OCS burst, ER visit, hospital, office visit); OR
- Member has Oral Corticosteroid dependent asthma; OR
- Member has BOTH:
  - A baseline Forced Expiratory Volume (FEV1) that is less than 80% predicted for adults and less than 90% for adolescents.
  - Prior drug therapy of either a leukotriene modifier OR med-high or max-tolerated ICS + controller OR max-tolerated ICS/LABA combo

#### **Chronic rhinosinusitis with nasal polyps (CRSwNP)**

Authorization of 6 months may be granted for members 12 years of age or older who have previously received a biologic drug indicated for CRSwNP in the past year.

OR

Authorization of 6 months may be granted for treatment of CRSwNP in members 12 years of age or older when all of the following criteria are met:

- Member has a confirmed diagnosis of CRSwNP
- The member has CRSwNP despite nasal surgery
- CRSwNP is inadequately controlled by medical therapy with 2 of the following in the past year, unless contraindicated or intolerant to:
  - Intranasal corticosteroids
  - Systemic corticosteroid therapy
  - Nasal budesonide nebulized solution

#### **Eosinophilic esophagitis (EoE)**

Authorization of 6 months may be granted for treatment of EoE in members 1 year of age or older, weighing at least 15 kg, when ALL of the following criteria are met:

- Diagnosis has been confirmed by esophageal biopsy as characterized by 15 or more intraepithelial esophageal eosinophils per high power field
- Member had a failure, intolerance, or contraindication to BOTH of the following:
  - Eight weeks of use of a generic proton pump inhibitor (e.g. esomeprazole, lansoprazole, omeprazole, pantoprazole, or rabeprazole)
  - Topical glucocorticoids (fluticasone using MDI without a spacer or budesonide administered as an oral slurry)

#### **Prurigo Nodularis**

Authorization of 6 months may be granted for members 18 years of age or older who have previously received a biologic drug indicated for prurigo nodularis in the past year.

OR

Authorization of 6 months may be granted for treatment of prurigo nodularis in members 18 years of age or older when ALL of the following criteria are met:



AETNA BETTER HEALTH®  
Coverage Policy/Guideline

Name: Dupixent

Page: 6 of 16

Effective Date: 10/16/2025

Last Review Date: 9/9/2025

Applies to:  Illinois

- Member has pruritus lasting at least 6 weeks.
- Member has history or signs of repeated itch-scratch cycle (e.g., scratching, picking, or rubbing).
- Member has a minimum of 20 nodular lesions.
- Member meets either of the following:
  - Member has had an inadequate response to one of the following:
    - A medium to super-high potency topical corticosteroid (see Appendix A)
    - A topical calcineurin inhibitor
    - Phototherapy (e.g., UVB, PUVA)
    - Pharmacologic treatment with methotrexate or cyclosporine
  - Member has had an intolerance or a clinical reason to avoid either of the following:
    - Medium to super-high potency topical corticosteroid (see Appendix A) and topical calcineurin inhibitor
    - Pharmacologic treatment with methotrexate and cyclosporine (see Appendix B)

#### **Chronic obstructive pulmonary disease (COPD)**

Authorization of 12 months may be granted for members 18 years of age or older who have previously received a biologic drug indicated for COPD in the past year.

OR

Authorization of 12 months may be granted for treatment of COPD in members 18 years of age or older when ALL of the following criteria are met:

- Diagnosis has been confirmed by spirometry showing forced expiratory volume in one second (FEV<sub>1</sub>)/forced vital capacity (FVC) less than 0.7 post-bronchodilation.
- Member demonstrates classic signs or symptoms of COPD (e.g., dyspnea, wheezing, chest tightness, fatigue, activity limitation, cough with or without sputum production, chronic bronchitis).
- Member has an absolute blood eosinophil count of at least 300 cells per microliter prior to initiating therapy with the requested medication.
- Member has inadequately controlled COPD as demonstrated by experiencing either of the following in the last year:
  - At least two moderate exacerbations resulting in treatment with systemic glucocorticoids, antibiotics, or both.
  - One or more severe exacerbation(s) requiring hospitalization or an emergency medical care visit.
- Member meets EITHER of the following:
  - Member is currently receiving maintenance inhaled triple therapy (i.e., inhaled corticosteroid [ICS], long-acting muscarinic antagonist [LAMA], and long-acting beta<sub>2</sub>-agonist [LABA]).
  - Member is currently receiving a LAMA and LABA, and has a contraindication to ICS.



AETNA BETTER HEALTH®  
Coverage Policy/Guideline

Name: Dupixent

Page: 7 of 16

Effective Date: 10/16/2025

Last Review Date: 9/9/2025

Applies to:  Illinois

- Member will continue to use maintenance COPD treatments (e.g., ICS with LAMA and LABA, LAMA and LABA) in combination with the requested medication.

### **Chronic spontaneous urticaria (CSU)**

Authorization of 6 months may be granted for members 12 years of age or older who have previously received a biologic drug indicated for CSU in the past year.

OR

Authorization of 6 months may be granted for treatment of CSU in members 12 years of age or older when ALL of the following criteria are met:

- Member remains symptomatic despite treatment with up-dosing (in accordance with EAACI/GA2LEN/EuroGuiDerm/APAAACI guidelines) of a second-generation H1 antihistamine (e.g., cetirizine, fexofenadine, levocetirizine, loratadine) for at least 2 weeks.
- Member has been evaluated for other causes of wheals (hives) and/or angioedema, including bradykinin-related angioedema and interleukin-1-associated urticarial syndromes (auto-inflammatory disorders, urticarial vasculitis).
- Member has experienced a spontaneous onset of wheals (hives), angioedema, or both, for at least 6 weeks.

### **Immune checkpoint inhibitor-related toxicity**

Authorization of 12 months may be granted for treatment of immune checkpoint inhibitor-related toxicity when the requested medication will be used for severe (G3) pruritus if no response to gabapentinoids in one month.

OR

Authorization of 12 months may be granted for treatment of immune checkpoint inhibitor-related toxicity when the requested medication will be used as additional therapy for severe (G3) or life-threatening (G4) bullous dermatitis.

### **Bullous Pemphigoid**

Authorization of 12 months may be granted for treatment of bullous pemphigoid in members 18 years of age or older when ALL of the following criteria are met:

- Diagnosis has been confirmed by EITHER of the following:
  - Direct immunofluorescence (DIF) study
  - Immune serological test(s) (e.g., Indirect immunofluorescence microscopy [IIF], ELISA)
- Member demonstrates characteristic clinical features of bullous pemphigoid (e.g., urticarial or eczematous or erythematous plaques, bullae, pruritus).
- Member has moderate to severe disease.
- Member meets EITHER of the following:
  - Member has had an inadequate treatment response with EITHER of the following:
    - A super-high potency topical corticosteroid (see Appendix A)



AETNA BETTER HEALTH®  
Coverage Policy/Guideline

Name:	Dupixent	Page:	8 of 16
Effective Date:	10/16/2025	Last Review Date:	9/9/2025
Applies to:	<input checked="" type="checkbox"/> Illinois		

- An oral corticosteroid
  - The use of super-high potency topical corticosteroid or oral corticosteroid is not advisable for the member (e.g., contraindications, prior intolerances).

## **Continuation of Therapy**

### **Atopic dermatitis**

Authorization of 12 months may be granted for members 6 months of age or older (including new members) who are using the requested medication for moderate-to-severe atopic dermatitis when the member has achieved or maintained a positive clinical response as evidenced by low disease activity (i.e., clear or almost clear skin), or improvement in signs and symptoms of atopic dermatitis (e.g., redness, itching, oozing/crusting).

### **Asthma**

Authorization of 12 months may be granted for continuation of treatment of moderate-to-severe asthma in members 6 years of age or older when BOTH of the following criteria are met:

- Asthma control has improved on the requested medication as demonstrated by at least ONE of the following:
  - A reduction in the frequency or severity of symptoms and exacerbations
  - A reduction in the daily maintenance oral corticosteroid dose
- Member will continue to use maintenance asthma treatments (e.g., inhaled corticosteroid, additional controller) in combination with the requested medication.

### **Chronic rhinosinusitis with nasal polyps (CRSwNP)**

Authorization of 12 months may be granted for continuation of treatment of CRSwNP in members 12 years of age or older when BOTH of the following are met:

- Member has achieved or maintained a positive clinical response with the requested medication as evidenced by improvement in signs and symptoms of CRSwNP (e.g., improvement in nasal congestion, nasal polyp size, loss of smell, anterior or posterior rhinorrhea, sino-nasal inflammation, hyposmia or facial pressure or pain, or reduction in corticosteroid use).

### **Eosinophilic Esophagitis (EoE)**

Authorization of 12 months may be granted for continuation of treatment of EoE in members 1 year of age or older, weighing at least 15 kg, when member has achieved or maintained a positive clinical response with the requested medication as evidenced by improvement in signs and symptoms of EoE (e.g., dysphagia, heartburn, chest pain, emesis).

### **Prurigo Nodularis**

Authorization of 12 months may be granted for members 18 years of age or older (including new members) who are using the requested medication for prurigo nodularis when the member has achieved or maintained a positive clinical response as evidenced by EITHER of the following:

- Low disease activity (i.e., clear or almost clear skin)



AETNA BETTER HEALTH®  
Coverage Policy/Guideline

Name: Dupixent

Page: 9 of 16

Effective Date: 10/16/2025

Last Review Date: 9/9/2025

Applies to:  Illinois

- Reduction in pruritic intensity and improvement in extent and severity of nodular lesions

### **Chronic obstructive pulmonary disease (COPD)**

Authorization of 12 months may be granted for continuation of treatment of COPD in members 18 years of age or older when BOTH of the following criteria are met:

- Member has achieved or maintained a positive clinical response as evidenced by improvement in signs and symptoms of COPD (e.g., decrease in exacerbations, improvement in pre-bronchodilator FEV<sub>1</sub>) or stabilization of disease.
- Member will continue to use maintenance COPD treatments (e.g., ICS with LAMA and LABA, LAMA and LABA) in combination with the requested medication.

### **Chronic spontaneous urticaria (CSU)**

Authorization of 12 months may be granted for continuation of treatment of chronic spontaneous urticaria in members 12 years of age or older, when the member has experienced a positive clinical response (e.g., improved symptoms, decrease in weekly urticaria activity score [UAS7]) since initiation of therapy.

### **Immune Checkpoint Inhibitor-Related Toxicities**

Authorization of 12 months may be granted for all members (including new members) who are using the requested medication for immune checkpoint inhibitor-related severe (G3) pruritus, severe (G3) or life-threatening (G4) bullous dermatitis, and who achieve or maintain a positive clinical response as evidenced by low disease activity or improvement in signs and symptoms of the condition..

### **Bullous Pemphigoid**

Authorization of 12 months may be granted for continuation of treatment of bullous pemphigoid in members 18 years of age or older, when the member has achieved or maintained a positive clinical response as evidenced by EITHER of the following:

- Low disease activity (e.g., absence of new or established lesions)
- Reduction in pruritus intensity and improvement in extent and severity of lesions

### **Other**

For all indications: Member cannot use the requested medication concomitantly with any other biologic drug or targeted synthetic drug for the same indication.

Note: If the member is a current smoker or vaper, they should be counseled on the harmful effects of smoking and vaping on pulmonary conditions and available smoking and vaping cessation options.

### **Appendix**

Appendix A: Table. Relative potency of select topical corticosteroid products



AETNA BETTER HEALTH®  
Coverage Policy/Guideline

Name: Dupixent Page: 10 of 16

Effective Date: 10/16/2025 Last Review Date: 9/9/2025

Applies to:  Illinois

Potency	Drug	Dosage form	Strength
Super-high potency (Group 1)	Augmented betamethasone dipropionate	Ointment, Lotion, Gel	0.05%
Super-high potency (Group 1)	Clobetasol propionate	Cream, Gel, Ointment, Solution, Cream (emollient), Lotion, Shampoo, Foam, Spray	0.05%
Super-high potency (Group 1)	Fluocinonide	Cream	0.1%
Super-high potency (Group 1)	Flurandrenolide	Tape	4 mcg/cm <sup>2</sup>
Super-high potency (Group 1)	Halobetasol propionate	Cream, Lotion, Ointment, Foam	0.05%
High potency (Group 2)	Amcinonide	Ointment	0.1%
High potency (Group 2)	Augmented betamethasone dipropionate	Cream	0.05%
High potency (Group 2)	Betamethasone dipropionate	Ointment	0.05%
High potency (Group 2)	Clobetasol propionate	Cream	0.025%
High potency (Group 2)	Desoximetasone	Cream, Ointment, Spray	0.25%
High potency (Group 2)	Desoximetasone	Gel	0.05%
High potency (Group 2)	Diflorasone diacetate	Ointment, Cream (emollient)	0.05%
High potency (Group 2)	Fluocinonide	Cream, Ointment, Gel, Solution	0.05%
High potency (Group 2)	Halcinonide	Cream, Ointment	0.1%
High potency (Group 2)	Halobetasol propionate	Lotion	0.01%
High potency (Group 3)	Amcinonide	Cream, Lotion	0.1%



AETNA BETTER HEALTH®  
Coverage Policy/Guideline

Name: Dupixent

Page: 11 of 16

Effective Date: 10/16/2025

Last Review Date: 9/9/2025

Applies to:  Illinois

Potency	Drug	Dosage form	Strength
High potency (Group 3)	Betamethasone dipropionate	Cream, hydrophilic emollient	0.05%
High potency (Group 3)	Betamethasone valerate	Ointment	0.1%
High potency (Group 3)	Betamethasone valerate	Foam	0.12%
High potency (Group 3)	Desoximetasone	Cream, Ointment	0.05%
High potency (Group 3)	Diflorasone diacetate	Cream	0.05%
High potency (Group 3)	Fluocinonide	Cream, aqueous emollient	0.05%
High potency (Group 3)	Fluticasone propionate	Ointment	0.005%
High potency (Group 3)	Mometasone furoate	Ointment	0.1%
High potency (Group 3)	Triamcinolone acetonide	Cream, Ointment	0.5%
Medium potency (Group 4)	Betamethasone dipropionate	Spray	0.05%
Medium potency (Group 4)	Clocortolone pivalate	Cream	0.1%
Medium potency (Group 4)	Fluocinolone acetonide	Ointment	0.025%
Medium potency (Group 4)	Flurandrenolide	Ointment	0.05%
Medium potency (Group 4)	Hydrocortisone valerate	Ointment	0.2%
Medium potency (Group 4)	Mometasone furoate	Cream, Lotion, Solution	0.1%
Medium potency (Group 4)	Triamcinolone acetonide	Cream	0.1%



AETNA BETTER HEALTH®  
Coverage Policy/Guideline

Name: Dupixent

Page: 12 of 16

Effective Date: 10/16/2025

Last Review Date: 9/9/2025

Applies to:  Illinois

Potency	Drug	Dosage form	Strength
Medium potency (Group 4)	Triamcinolone acetonide	Ointment	0.05% and 0.1%
Medium potency (Group 4)	Triamcinolone acetonide	Aerosol Spray	0.2 mg per 2-second spray
Lower-mid potency (Group 5)	Betamethasone dipropionate	Lotion	0.05%
Lower-mid potency (Group 5)	Betamethasone valerate	Cream	0.1%
Lower-mid potency (Group 5)	Desonide	Ointment, Gel	0.05%
Lower-mid potency (Group 5)	Fluocinolone acetonide	Cream	0.025%
Lower-mid potency (Group 5)	Flurandrenolide	Cream, Lotion	0.05%
Lower-mid potency (Group 5)	Fluticasone propionate	Cream, Lotion	0.05%
Lower-mid potency (Group 5)	Hydrocortisone butyrate	Cream, Lotion, Ointment, Solution	0.1%
Lower-mid potency (Group 5)	Hydrocortisone probutate	Cream	0.1%
Lower-mid potency (Group 5)	Hydrocortisone valerate	Cream	0.2%
Lower-mid potency (Group 5)	Prednicarbate	Cream (emollient), Ointment	0.1%
Lower-mid potency (Group 5)	Triamcinolone acetonide	Lotion	0.1%



AETNA BETTER HEALTH®  
Coverage Policy/Guideline

Name: Dupixent

Page: 13 of 16

Effective Date: 10/16/2025

Last Review Date: 9/9/2025

Applies to:  Illinois

Potency	Drug	Dosage form	Strength
Lower-mid potency (Group 5)	Triamcinolone acetonide	Ointment	0.025%
Low potency (Group 6)	Alclometasone dipropionate	Cream, Ointment	0.05%
Low potency (Group 6)	Betamethasone valerate	Lotion	0.1%
Low potency (Group 6)	Desonide	Cream, Lotion, Foam	0.05%
Low potency (Group 6)	Fluocinolone acetonide	Cream, Solution, Shampoo, Oil	0.01%
Low potency (Group 6)	Triamcinolone acetonide	Cream, lotion	0.025%
Least potent (Group 7)	Hydrocortisone (base, greater than or equal to 2%)	Cream, Ointment, Solution	2.5%
Least potent (Group 7)	Hydrocortisone (base, greater than or equal to 2%)	Lotion	2%
Least potent (Group 7)	Hydrocortisone (base, less than 2%)	Cream, Ointment, Gel, Lotion, Spray, Solution	1%
Least potent (Group 7)	Hydrocortisone (base, less than 2%)	Cream, Ointment	0.5%
Least potent (Group 7)	Hydrocortisone acetate	Cream	2.5%
Least potent (Group 7)	Hydrocortisone acetate	Lotion	2%
Least potent (Group 7)	Hydrocortisone acetate	Cream	1%

#### Appendix B: Examples of Clinical Reasons to Avoid Pharmacologic Treatment with Methotrexate or Cyclosporine

- Clinical diagnosis of alcohol use disorder, alcoholic liver disease or other chronic liver disease
- Drug interaction
- Risk of treatment-related toxicity
- Pregnancy or currently planning pregnancy
- Breastfeeding
- Significant comorbidity prohibits use of systemic agents (e.g., liver or kidney disease, blood dyscrasias, uncontrolled hypertension)



AETNA BETTER HEALTH®  
Coverage Policy/Guideline

Name:	Dupixent	Page:	14 of 16
Effective Date:	10/16/2025	Last Review Date:	9/9/2025
Applies to:	<input checked="" type="checkbox"/> Illinois		

- Hypersensitivity
- History of intolerance or adverse event

### Approval Duration and Quantity Restrictions:

#### Approval:

- Initial: 6 months
- Renewal: 12 months

#### Quantity Level Limit:

- Dupixent 200 mg/ 1.14 mL pre-filled syringe/pen: 2 syringes/pens per 28 days
- Dupixent 300 mg/ 2 mL pre-filled syringe/pen: 4 syringes/pens per 28 days
- Dupixent 100 mg/ 0.67 mL pre-filled syringe: 2 syringes per 28 days

NOTE: Quantity approved with requests will be based upon FDA-approved dosage.

### References:

1. Dupixent [package insert]. Tarrytown, NY: Regeneron Pharmaceuticals, Inc.; June 2025.
2. Sidbury R, Alikhan A, Bercovitch L, et al. Guidelines of care for the management of atopic dermatitis in adults with topical therapies. *J Am Acad Dermatol.* 2023;89(1):e1-e20.
3. Simpson EL, Bieber T, Guttman-Yassky E, et al. Two phase 3 trials of dupilumab versus placebo in atopic dermatitis. *N Engl J Med.* 2016;375:2335-2348.
4. Castro M, Corren J, Pavord ID, et al. Dupilumab Efficacy and Safety in Moderate-to-Severe Uncontrolled Asthma. *N Engl J Med.* 2018;378(26):2486-2496.
5. Rabe KF, Nair P, Brusselle G, et al. Efficacy and Safety of Dupilumab in Glucocorticoid-Dependent Severe Asthma. *N Engl J Med.* 2018;378(26):2475-2485.
6. Global Initiative for Asthma (GINA). Global Strategy for Asthma Management and Prevention. 2024 update. Available at: [https://ginasthma.org/wp-content/uploads/2024/05/GINA-2024-Strategy-Report-24\\_05\\_22\\_WMS.pdf](https://ginasthma.org/wp-content/uploads/2024/05/GINA-2024-Strategy-Report-24_05_22_WMS.pdf). Accessed March 1, 2025.
7. Topical Corticosteroids. Drug Facts and Comparisons. Facts & Comparisons [database online]. St. Louis, MO: Wolters Kluwer Health Inc; July 18, 2024. Accessed November 9, 2024.
8. ClinicalTrials.gov. National Library of Medicine (US). Identifier NCT02912468. A Controlled Clinical Study of Dupilumab in Patients with Nasal Polyps (SINUS-24) 2016 Sep 23. Available from: <https://clinicaltrials.gov/ct2/show/NCT02912468>.
9. ClinicalTrials.gov. National Library of Medicine (US). Identifier NCT02898454. A Controlled Clinical Study of Dupilumab in Patients with Nasal Polyps (SINUS-52) 2016 Sep 13. Available from: <https://clinicaltrials.gov/ct2/show/NCT02898454>.
10. Fishbein AB, Silverberg JI, Wilson EJ, et al. Update on atopic dermatitis: Diagnosis, severity assessment, and treatment selection. *J Allergy Clin Immunol Pract.* 2020;8(1): 91-101.
11. Cloutier MM, Dixon AE, Krishnan JA, et al. Managing asthma in adolescents and adults: 2020 asthma guideline update from the National Asthma Education and Prevention Program. *JAMA.* 2020;324(22): 2301-2317.
12. Bachert C, Han JK, Wagenmann M, et al. EUFOREA expert board meeting on uncontrolled severe chronic rhinosinusitis with nasal polyps (CRSwNP) and biologics: Definitions and management. *J Allergy Clin Immunol.* 2021;147(1):29-36.



AETNA BETTER HEALTH®  
Coverage Policy/Guideline

Name:	Dupixent	Page:	15 of 16
Effective Date:	10/16/2025	Last Review Date:	9/9/2025
Applies to:	<input checked="" type="checkbox"/> Illinois		

13. Lucendo AJ, Molina-Infante J, Arias A, et al. Guidelines on eosinophilic esophagitis: evidence-based statements and recommendations for diagnosis and management in children and adults. *United European Gastroenterol J.* 2017;5(3):355-358.
14. Gonsalves NP, Aceves S. Diagnosis and treatment of eosinophilic esophagitis. *J Allergy Clin Immunol.* 2020;145(1):1-7.
15. ClinicalTrials.gov. National Library of Medicine (US). Identifier NCT03633617. Study to determine the efficacy and safety of Dupilumab in Adult and Adolescent Patients with Eosinophilic Esophagitis (EoE) 2022 May 27. Available from: <https://clinicaltrials.gov/ct2/show/NCT03633617>.
16. ClinicalTrials.gov. National Library of Medicine (US). Identifier NCT03346434. Safety, Pharmacokinetics and Efficacy of Dupilumab in Patients ≥6 months to <6 years with Moderate-to-Severe Atopic Dermatitis (Liberty AD PRESCHOOL) 2022 Jun 10. Available from: <https://clinicaltrials.gov/ct2/show/NCT03346434>.
17. Fokkens WJ, Lund VJ, Hopkins C, et al. European Position Paper on Rhinosinusitis and Nasal Polyps 2020. *Rhinology.* 2020;58(Suppl S29):1-464.
18. Hopkins C. Chronic Rhinosinusitis with Nasal Polyps. *N Engl J Med.* 2019;381(1):55-63.
19. ClinicalTrials.gov. National Library of Medicine (US). Identifier NCT04183335. Study of Dupilumab for the Treatment of Patients With Prurigo Nodularis, Inadequately Controlled on Topical Prescription Therapies or When Those Therapies Are Not Advisable (LIBERTY-PN PRIME). 2022 February 17. Available from: <https://clinicaltrials.gov/ct2/show/NCT04183335>.
20. ClinicalTrials.gov. National Library of Medicine (US). Identifier NCT04202679. Study of Dupilumab for the Treatment of Patients With Prurigo Nodularis, Inadequately Controlled on Topical Prescription Therapies or When Those Therapies Are Not Advisable (PRIME2). 2022 September 28. Available from: <https://clinicaltrials.gov/ct2/show/NCT04202679>.
21. Ständer HF, Elmariah S, Zeidler C, et al. Diagnostic and treatment algorithm for chronic nodular prurigo. *J Am Acad Dermatol.* 2020;82(2):460-468.
22. Elmariah S, Kim B, Berger T, et al. Practical approaches for diagnosis and management of prurigo nodularis: United States expert panel consensus. *J Am Acad Dermatol.* 2021;84(3):747-760.
23. Cyclosporine. Drug Facts and Comparisons. Facts & Comparisons [database online]. St. Louis, MO: Wolters Kluwer Health Inc; November 6, 2024. Accessed November 10, 2024.
24. Methotrexate. Drug Facts and Comparisons. Facts & Comparisons [database online]. St. Louis, MO: Wolters Kluwer Health Inc; November 7, 2024. Accessed November 10, 2024.
25. The NCCN Drugs & Biologics Compendium® © 2025 National Comprehensive Cancer Network, Inc. Available at: <http://www.nccn.org>. Accessed March 3, 2025.
26. NCCN Clinical Practice Guidelines in Oncology® (NCCN Guidelines®). Management of Immune Checkpoint-Related Toxicities. Version 1.2025. Available at: [www.nccn.org](http://www.nccn.org). Accessed March 3, 2025.
27. ClinicalTrials.gov. National Library of Medicine (US). Identifier NCT04394351. Study to Investigate the Efficacy and Safety of Dupilumab in Pediatric Patients With Active Eosinophilic Esophagitis (EoE) (EoE KIDS). 2023 June 05. Available from: <https://clinicaltrials.gov/ct2/show/NCT04394351>.
28. Lucendo AJ, Sánchez-Cazalilla M. Adult versus pediatric eosinophilic esophagitis: important differences and similarities for the clinician to understand. *Expert Rev Clin Immunol.* 2012;8(8):733-45.



AETNA BETTER HEALTH®  
Coverage Policy/Guideline

Name: Dupixent Page: 16 of 16

Effective Date: 10/16/2025 Last Review Date: 9/9/2025

Applies to:  Illinois

29. Global Initiative for Chronic Obstructive Lung Disease (GOLD). Global Strategy for the Diagnosis, Management, and Prevention of Chronic Obstructive Pulmonary Disease (2024 Report). Available at: <https://goldcopd.org/2024-gold-report/>. Accessed March 2025.
30. Bhatt SP, Rabe KF, Hanania NA, et al. Dupilumab for COPD with Type 2 Inflammation Indicated by Eosinophil Counts. *N Engl J Med*. 2023;389(3):205-214.
31. ClinicalTrials.gov. National Library of Medicine (US). Identifier NCT04456673. Pivotal Study to Assess the Efficacy, Safety and Tolerability of Dupilumab in Patients with Moderate to Severe COPD with Type 2 Inflammation (NOTUS). Last updated October 15, 2024. Accessed 2025 March 12. Available from: <https://clinicaltrials.gov/ct2/show/NCT04456673>.
32. Dellon E, Muir AB, Katzka DA, et al. ACG Clinical Guideline: Diagnosis and Management of Eosinophilic Esophagitis. *AJC*. 2025;120(1):p31-59.
33. Rank MA, Chu DK, Bognanni A, et al. The Joint Task Force on Practice Parameters GRADE guidelines for the medical management of chronic rhinosinusitis with nasal polyposis. *J Allergy Clin Immunol*. 2023 Feb;151(2):386-398.
34. Zuberbier T, Abdul Latiff AH, Abuzakouk M, et al. The international EAACI/GA<sup>2</sup>LEN/EuroGuiDerm/APAAACI guideline for the definition, classification, diagnosis, and management of urticaria. *Allergy*. 2022 Mar;77(3):734-766.
35. Bernstein DI, Blessing-Moore J, Cox L, et al. The diagnosis and management of acute and chronic urticaria: 2014 update. American Academy of Allergy, Asthma & Immunology Practice Parameter. <http://www.aaaai.org/practice-resources/statements-and-practice-parameters/practice-parameter-guidelines.aspx>. Accessed April 23, 2025.
36. Borradori L, Van Beek N, Feliciani C, et al. Updated S2 K guidelines for the management of bullous pemphigoid initiated by the European Academy of Dermatology and Venereology (EADV). *J Eur Acad Dermatol Venereol*. 2022 Oct;36(10):1689-1704.
37. ClinicalTrials.gov. National Library of Medicine (US). Identifier NCT04206553. A Study to Evaluate the Efficacy and Safety of Dupilumab in Adult Patients With Bullous Pemphigoid (LIBERTY-BP). Last updated February 6, 2025. Accessed 2025 July 1. Available from: <https://clinicaltrials.gov/study/NCT04206553>.
38. Murrell DF, Joly P, Werth VP, et al. Study Design of a Phase 2/3 Randomized Controlled Trial of Dupilumab in Adults with Bullous Pemphigoid: LIBERTY-BP ADEPT. *Adv Ther*. 2024 Jul;41(7):2991-3002.