



AETNA BETTER HEALTH®  
Coverage Policy/Guideline

Name: Enbrel Page: 1 of 14

Effective Date: 2/11/2026 Last Review Date: 1/2026

Applies to:  Illinois  Florida  Florida Kids  
 New Jersey  Maryland  Michigan  
 Pennsylvania Kids  Virginia  Kentucky PRMD

### Intent:

The intent of this policy/guideline is to provide information to the prescribing practitioner outlining the coverage criteria for Enbrel under the patient's prescription drug benefit.

### Description:

The indications below including FDA-approved indications and compendial uses are considered a covered benefit provided that all the approval criteria are met and the member has no exclusions to the prescribed therapy.

#### FDA-approved Indications<sup>1</sup>

- Moderately to severely active rheumatoid arthritis (RA)
- Moderately to severely active polyarticular juvenile idiopathic arthritis (pJIA) in patients 2 years of age and older
- Active psoriatic arthritis (PsA)
- Active ankylosing spondylitis (AS)
- Chronic moderate to severe plaque psoriasis (PsO) in patients 4 years or older who are candidates for systemic therapy or phototherapy - Reference the Biological Response Modifiers (BRMs) in the Treatment of Plaque Psoriasis NJ Protocol
- Active juvenile psoriatic arthritis (JPsA) in pediatric patients 2 years of age and older

#### Compendial Uses

- Non-radiographic axial spondyloarthritis<sup>2,23</sup>
- Oligoarticular juvenile idiopathic arthritis<sup>26</sup>
- Reactive arthritis<sup>4</sup>
- Hidradenitis suppurativa, severe, refractory<sup>4</sup>
- Behcet's disease<sup>4,17</sup>
- Graft versus host disease<sup>4,16,22,23</sup>
- Immune checkpoint inhibitor-related toxicity<sup>22</sup>

All other indications are considered experimental/investigational and not medically necessary.

### Applicable Drug List:

#### Preferred Agent:

Enbrel



AETNA BETTER HEALTH®  
Coverage Policy/Guideline

Name: Enbrel Page: 2 of 14

Effective Date: 2/11/2026 Last Review Date: 1/2026

Applies to:  Illinois  Florida  Florida Kids  
 New Jersey  Maryland  Michigan  
 Pennsylvania Kids  Virginia  Kentucky PRMD

## Policy/Guideline:

### Documentation

Submission of the following information is necessary to initiate the prior authorization review:

#### Rheumatoid arthritis (RA)

##### Initial requests

- Chart notes, medical record documentation, or claims history supporting previous medications tried (if applicable), including response to therapy. If therapy is not advisable, documentation of clinical reason to avoid therapy.
- Laboratory results, chart notes, or medical record documentation of biomarker testing (i.e., rheumatoid factor [RF], anti-cyclic citrullinated peptide [anti-CCP], and C-reactive protein [CRP] and/or erythrocyte sedimentation rate [ESR]) (if applicable).

##### Continuation requests

Chart notes or medical record documentation supporting positive clinical response.

#### Articular juvenile idiopathic arthritis (JIA)

##### Initial requests

Chart notes, medical record documentation, or claims history supporting previous medications tried (if applicable), including response to therapy.

##### Continuation requests

Chart notes or medical record documentation supporting positive clinical response.

#### Psoriatic arthritis (PsA), ankylosing spondylitis (AS), non-radiographic axial spondyloarthritis (nr-axSpA), reactive arthritis, hidradenitis suppurativa, immune checkpoint inhibitor-related inflammatory arthritis, and chronic graft versus host disease

##### Initial requests

Chart notes, medical record documentation, or claims history supporting previous medications tried (if applicable), including response to therapy. If therapy is not advisable, documentation of clinical reason to avoid therapy.

##### Continuation requests

Chart notes or medical record documentation supporting positive clinical response.



AETNA BETTER HEALTH®  
Coverage Policy/Guideline

Name: Enbrel Page: 3 of 14

Effective Date: 2/11/2026 Last Review Date: 1/2026

Applies to:  Illinois  Florida  Florida Kids  
 New Jersey  Maryland  Michigan  
 Pennsylvania Kids  Virginia  Kentucky PRMD

Acute graft versus host disease and immune checkpoint inhibitor-related toxicity (initial requests only)

Chart notes, medical record documentation, or claims history supporting previous medications tried (if applicable), including response to therapy. If therapy is not advisable, documentation of clinical reason to avoid therapy.

Behcet's disease (initial requests only)

Chart notes, medical record documentation, or claims history supporting previous medications tried, including response to therapy (if applicable).

**Prescriber Specialties**

This medication must be prescribed by or in consultation with one of the following:

- Rheumatoid arthritis, articular juvenile idiopathic arthritis, ankylosing spondylitis, non-radiographic axial spondyloarthritis, reactive arthritis, and Behcet's disease: rheumatologist
- Psoriatic arthritis and hidradenitis suppurativa: rheumatologist or dermatologist
- Graft versus host disease: oncologist or hematologist
- Immune checkpoint inhibitor-related inflammatory arthritis: oncologist, hematologist, or rheumatologist
- Immune checkpoint inhibitor-related toxicity: oncologist, hematologist, or dermatologist

**Coverage Criteria**

Rheumatoid arthritis (RA)<sup>1,5-7,30,31</sup>

Authorization of 12 months may be granted for adult members who have previously received a biologic or targeted synthetic drug indicated for moderately to severely active rheumatoid arthritis (RA) within the past 120 days.

Authorization of 12 months may be granted for adult members for treatment of moderately to severely active RA when both of the following criteria are met:

- Member meets either of the following:
  - Member has been tested for either of the following biomarkers and the test was positive:
    - Rheumatoid factor (RF)



AETNA BETTER HEALTH®  
Coverage Policy/Guideline

Name: Enbrel Page: 4 of 14

Effective Date: 2/11/2026 Last Review Date: 1/2026

Applies to:

<input type="checkbox"/> Illinois	<input type="checkbox"/> Florida	<input type="checkbox"/> Florida Kids
<input checked="" type="checkbox"/> New Jersey	<input type="checkbox"/> Maryland	<input type="checkbox"/> Michigan
<input type="checkbox"/> Pennsylvania Kids	<input type="checkbox"/> Virginia	<input type="checkbox"/> Kentucky PRMD

- Anti-cyclic citrullinated peptide (anti-CCP)
- Member has been tested for ALL of the following biomarkers:
  - RF
  - Anti-CCP
  - C-reactive protein (CRP) and/or erythrocyte sedimentation rate (ESR)
- Member meets ONE of the following:
  - Member has failed to achieve a low disease activity after a 3-month trial of methotrexate (MTX) monotherapy at a maximum titrated dose of at least 15 mg per week and meets any of the following conditions:
    - Member has had a documented inadequate response to MTX in combination with at least one other conventional synthetic drug (i.e., hydroxychloroquine and/or sulfasalazine) after a 3-month trial at a maximum tolerated dose(s).
    - Member has experienced a documented intolerable adverse event to hydroxychloroquine or sulfasalazine.
    - Member has a documented contraindication to hydroxychloroquine (see Appendix A) and sulfasalazine (e.g., porphyria, intestinal or urinary obstruction).
    - Member has moderate to high disease activity.
  - Member was unable to tolerate a 3-month trial of MTX monotherapy at a maximum titrated dose of at least 15 mg per week and meets any of the following conditions:
    - Member has had a documented inadequate response to MTX in combination with at least one other conventional synthetic drug (i.e., hydroxychloroquine and/or sulfasalazine) after a 3-month trial at a maximum tolerated dose(s).
    - Member has stopped taking MTX and has had a documented inadequate response to another conventional synthetic drug (i.e., leflunomide, hydroxychloroquine, and/or sulfasalazine) alone or in combination after a 3-month trial at a maximum tolerated dose(s).
    - Member has experienced a documented intolerable adverse event to leflunomide, hydroxychloroquine, or sulfasalazine.



AETNA BETTER HEALTH®  
Coverage Policy/Guideline

Name: Enbrel Page: 5 of 14

Effective Date: 2/11/2026 Last Review Date: 1/2026

Applies to:

<input type="checkbox"/> Illinois	<input type="checkbox"/> Florida	<input type="checkbox"/> Florida Kids
<input checked="" type="checkbox"/> New Jersey	<input type="checkbox"/> Maryland	<input type="checkbox"/> Michigan
<input type="checkbox"/> Pennsylvania Kids	<input type="checkbox"/> Virginia	<input type="checkbox"/> Kentucky PRMD

- Member has a documented contraindication to leflunomide, hydroxychloroquine (see Appendix A), and sulfasalazine (e.g., porphyria, intestinal or urinary obstruction).
- Member has moderate to high disease activity.
- Member has experienced a documented intolerable adverse event or has a documented contraindication to MTX (see Appendix A), discontinues MTX, and meets any of the following conditions:
  - Member has had a documented inadequate response to another conventional synthetic drug (i.e., leflunomide, hydroxychloroquine, and/or sulfasalazine) alone or in combination after a 3-month trial at a maximum tolerated dose(s).
  - Member has experienced a documented intolerable adverse event to leflunomide, hydroxychloroquine, or sulfasalazine.
  - Member has a documented contraindication to leflunomide, hydroxychloroquine, (see Appendix A), and sulfasalazine (e.g., porphyria, intestinal or urinary obstruction).
  - Member has moderate to high disease activity.

#### Articular juvenile idiopathic arthritis (JIA)<sup>1,8,26</sup>

Authorization of 12 months may be granted for members 2 years of age or older who have previously received a biologic or targeted synthetic drug indicated for moderately to severely active articular juvenile idiopathic arthritis.

Authorization of 12 months may be granted for members 2 years of age or older for treatment of moderately to severely active articular juvenile idiopathic arthritis when any of the following criteria is met:

- Member has had an inadequate response to methotrexate or another conventional synthetic drug (e.g., leflunomide, sulfasalazine, hydroxychloroquine) administered at an adequate dose and duration.
- Member has had an inadequate response to a trial of scheduled non-steroidal anti-inflammatory drugs (NSAIDs) and/or intra-articular glucocorticoids (e.g., triamcinolone hexacetonide) and one of the following risk factors for poor outcome:
  - Involvement of ankle, wrist, hip, sacroiliac joint, and/or temporomandibular joint (TMJ)
  - Presence of erosive disease or enthesitis
  - Delay in diagnosis



AETNA BETTER HEALTH®  
Coverage Policy/Guideline

Name: Enbrel Page: 6 of 14

Effective Date: 2/11/2026 Last Review Date: 1/2026

Applies to:  Illinois  Florida  Florida Kids  
 New Jersey  Maryland  Michigan  
 Pennsylvania Kids  Virginia  Kentucky PRMD

- Elevated levels of inflammation markers
- Symmetric disease
- Member has risk factors for disease severity and potentially a more refractory disease course (see Appendix B) and meets one of the following:
  - High-risk joints are involved (e.g., cervical spine, wrist, or hip)
  - High disease activity
  - Is judged to be at high risk for disabling joint disease

#### Psoriatic arthritis (PsA)<sup>1,10-13,20</sup>

Authorization of 12 months may be granted for members 2 years of age or older who have previously received a biologic or targeted synthetic drug indicated for active psoriatic arthritis.

Authorization of 12 months may be granted for members 2 years of age or older for treatment of active psoriatic arthritis when either of the following criteria is met:

- Member has mild to moderate disease and meets one of the following criteria:
  - Member has had an inadequate response to methotrexate, leflunomide, or another conventional synthetic drug (e.g., sulfasalazine) administered at an adequate dose and duration.
  - Member has an intolerance or contraindication to methotrexate or leflunomide (see Appendix A), or another conventional synthetic drug (e.g., sulfasalazine).
  - Member has enthesitis or predominantly axial disease.
- Member has severe disease.

#### Ankylosing spondylitis (AS) and non-radiographic axial spondyloarthritis (nr-axSpA)<sup>1,2,14,15</sup>

Authorization of 12 months may be granted for adult members who have previously received a biologic or targeted synthetic drug indicated for active ankylosing spondylitis or active non-radiographic axial spondyloarthritis.

Authorization of 12 months may be granted for adult members for treatment of active ankylosing spondylitis or active non-radiographic axial spondyloarthritis when any of the following criteria is met:

- Member has had an inadequate response to at least two non-steroidal anti-inflammatory drugs (NSAIDs).
- Member has an intolerance or contraindication to two or more NSAIDs.



AETNA BETTER HEALTH®  
Coverage Policy/Guideline

Name: Enbrel Page: 7 of 14

Effective Date: 2/11/2026 Last Review Date: 1/2026

Applies to:  Illinois  Florida  Florida Kids  
 New Jersey  Maryland  Michigan  
 Pennsylvania Kids  Virginia  Kentucky PRMD

#### Reactive arthritis<sup>3,4,21,31,32</sup>

Authorization of 12 months may be granted for members who have previously received a biologic indicated for reactive arthritis.

Authorization of 12 months may be granted for treatment of reactive arthritis when either of the following criteria is met:

- Member has had an inadequate response to methotrexate or sulfasalazine.
- Member has an intolerance or contraindication to methotrexate (see Appendix A) and sulfasalazine (e.g., porphyria, intestinal or urinary obstruction).

#### Hidradenitis suppurativa<sup>4,28,29</sup>

Authorization of 12 months may be granted for members who have previously received a biologic indicated for the treatment of severe, refractory hidradenitis suppurativa.

Authorization of 12 months may be granted for treatment of severe, refractory hidradenitis suppurativa when either of the following is met:

- Member has had an inadequate response to an oral antibiotic used for the treatment of hidradenitis suppurativa for at least 90 days (e.g., clindamycin, metronidazole, moxifloxacin, rifampin, tetracyclines).
- Member has an intolerance or contraindication to oral antibiotics used for the treatment of hidradenitis suppurativa.

#### Graft versus host disease<sup>4,16,22,23</sup>

Authorization of 12 months may be granted for treatment of graft versus host disease when either of the following criteria is met:

- Member has had an inadequate response to systemic corticosteroids.
- Member has an intolerance or contraindication to corticosteroids.

#### Behcet's disease<sup>4,17</sup>

Authorization of 12 months may be granted for members who have previously received Otezla or a biologic indicated for the treatment of Behcet's disease.

Authorization of 12 months may be granted for the treatment of Behcet's disease when the member has had an inadequate response to at least one non-biologic medication for Behcet's disease (e.g., azathioprine, colchicine, cyclosporine, systemic corticosteroids).



AETNA BETTER HEALTH®  
Coverage Policy/Guideline

Name: Enbrel Page: 8 of 14

Effective Date: 2/11/2026 Last Review Date: 1/2026

Applies to:

<input type="checkbox"/> Illinois	<input type="checkbox"/> Florida	<input type="checkbox"/> Florida Kids
<input checked="" type="checkbox"/> New Jersey	<input type="checkbox"/> Maryland	<input type="checkbox"/> Michigan
<input type="checkbox"/> Pennsylvania Kids	<input type="checkbox"/> Virginia	<input type="checkbox"/> Kentucky PRMD

### Immune checkpoint inhibitor-related toxicity<sup>22</sup>

Authorization of 1 month may be granted for treatment of immune checkpoint inhibitor-related toxicity when the member has Stevens-Johnson syndrome or toxic epidermal necrolysis.

Authorization of 12 months may be granted for treatment of immune checkpoint inhibitor-related toxicity when the member has moderate or severe immunotherapy-related inflammatory arthritis and either of the following is met:

- Member has had an inadequate response to corticosteroids or a conventional synthetic drug (e.g., methotrexate, sulfasalazine, leflunomide, hydroxychloroquine).
- Member has an intolerance or contraindication to corticosteroids and a conventional synthetic drug (e.g., methotrexate, sulfasalazine, leflunomide, hydroxychloroquine).

### **Continuation of Therapy**

#### Rheumatoid arthritis (RA)<sup>1,5-7,30,31</sup>

Authorization of 12 months may be granted for all adult members (including new members) who are using the requested medication for moderately to severely active rheumatoid arthritis and who achieve or maintain a positive clinical response as evidenced by disease activity improvement of at least 20% from baseline in tender joint count, swollen joint count, pain, or disability.

#### Articular juvenile idiopathic arthritis (JIA)<sup>1,8</sup>

Authorization of 12 months may be granted for all members 2 years of age or older (including new members) who are using the requested medication for moderately to severely active articular juvenile idiopathic arthritis and who achieve or maintain a positive clinical response as evidenced by low disease activity or improvement in signs and symptoms of the condition when there is improvement in any of the following from baseline:

- Number of joints with active arthritis (e.g., swelling, pain, limitation of motion)
- Number of joints with limitation of movement
- Functional ability

#### Psoriatic arthritis (PsA)<sup>1,10-13,20</sup>

Authorization of 12 months may be granted for all members 2 years of age or older (including new members) who are using the requested medication for psoriatic arthritis and who achieve or maintain a positive clinical response as evidenced by low disease activity or



AETNA BETTER HEALTH®  
Coverage Policy/Guideline

Name: Enbrel Page: 9 of 14

Effective Date: 2/11/2026 Last Review Date: 1/2026

Applies to:  Illinois  Florida  Florida Kids  
 New Jersey  Maryland  Michigan  
 Pennsylvania Kids  Virginia  Kentucky PRMD

improvement in signs and symptoms of the condition when there is improvement in any of the following from baseline:

- Number of swollen joints
- Number of tender joints
- Dactylitis
- Enthesitis
- Axial disease
- Skin and/or nail involvement
- Functional status
- C-reactive protein (CRP)

#### Ankylosing spondylitis (AS) and non-radiographic axial spondyloarthritis (nr-axSpA)<sup>1,2,14,15</sup>

Authorization of 12 months may be granted for all adult members (including new members) who are using the requested medication for ankylosing spondylitis or non-radiographic axial spondyloarthritis and who achieve or maintain a positive clinical response as evidenced by low disease activity or improvement in signs and symptoms of the condition when there is improvement in any of the following from baseline:

- Functional status
- Total spinal pain
- Inflammation (e.g., morning stiffness)
- Swollen joints
- Tender joints
- C-reactive protein (CRP)

#### Reactive arthritis<sup>3,21</sup>

Authorization of 12 months may be granted for all members (including new members) who are using the requested medication for reactive arthritis and who achieve or maintain a positive clinical response as evidenced by low disease activity or improvement in signs and symptoms of the condition (e.g., tender joint count, swollen joint count, or pain).

#### Hidradenitis suppurativa<sup>4,28,29</sup>

Authorization of 12 months may be granted for all members (including new members) who are using the requested medication for severe, refractory hidradenitis suppurativa and who achieve or maintain a positive clinical response as evidenced by low disease activity or improvement in signs and symptoms of the condition when any of the following is met:

- Reduction in abscess and inflammatory nodule count from baseline



AETNA BETTER HEALTH®  
Coverage Policy/Guideline

Name: Enbrel Page: 10 of 14

Effective Date: 2/11/2026 Last Review Date: 1/2026

Applies to:  Illinois  Florida  Florida Kids  
 New Jersey  Maryland  Michigan  
 Pennsylvania Kids  Virginia  Kentucky PRMD

- Reduced formation of new sinus tracts and scarring
- Decrease in frequency of inflammatory lesions from baseline
- Reduction in pain from baseline
- Reduction in suppuration from baseline
- Improvement in frequency of relapses from baseline
- Improvement in quality of life from baseline
- Improvement on a disease severity assessment tool from baseline

#### Acute graft versus host disease and immune checkpoint inhibitor-related toxicity

All members (including new members) requesting authorization for continuation of therapy must meet all requirements in the coverage criteria.

#### Chronic graft versus host disease

Authorization of 12 months may be granted for all members (including new members) who are using the requested medication for chronic graft versus host disease and who achieve or maintain a positive clinical response as evidenced by low disease activity or improvement in signs and symptoms of the condition.

#### Behcet's disease

Authorization of 12 months may be granted for all members (including new members) who are using the requested medication for Behcet's disease and who achieve or maintain a positive clinical response as evidenced by low disease activity or improvement in signs and symptoms of the condition.

#### Immune checkpoint inhibitor-related inflammatory arthritis

Authorization of 12 months may be granted for all members (including new members) who are using the requested medication for immunotherapy-related inflammatory arthritis and who achieve or maintain a positive clinical response as evidenced by low disease activity or improvement in signs and symptoms of the condition.

#### **Other<sup>1,19</sup>**

For all indications: Member has had a documented negative tuberculosis (TB) test (which can include a tuberculosis skin test [TST] or an interferon-release assay [IGRA] within 12 months of initiating therapy for persons who are naïve to biologic drugs or targeted synthetic drugs associated with an increased risk of TB.



AETNA BETTER HEALTH®  
Coverage Policy/Guideline

Name: Enbrel Page: 11 of 14

Effective Date: 2/11/2026 Last Review Date: 1/2026

Applies to:  Illinois  Florida  Florida Kids  
 New Jersey  Maryland  Michigan  
 Pennsylvania Kids  Virginia  Kentucky PRMD

If the screening testing for TB is positive, there must be further testing to confirm there is no active disease (e.g., chest x-ray). Do not administer the requested medication to members with active TB infection. If there is latent disease, TB treatment must be started before initiation of the requested medication.

For all indications: Member cannot use the requested medication concomitantly with any other biologic drug or targeted synthetic drug for the same indication.

### Dosage And Administration

Approvals may be subject to dosing limits in accordance with FDA-approved labeling, accepted compendia, and/or evidence-based practice guidelines. Dose optimization with 50 mg product formulations should be used when possible. Exceptions for higher quantities of 25 mg vials will be allowed when the member has a latex allergy or is following FDA-approved weight-based dosing.

### Appendix

Appendix A: Examples of Clinical Reasons to Avoid Pharmacologic Treatment with Methotrexate, Hydroxychloroquine, Leflunomide, Cyclosporine, or Acitretin<sup>25</sup>

- Clinical diagnosis of alcohol use disorder, alcoholic liver disease, or other chronic liver disease
- Drug interaction
- Risk of treatment-related toxicity
- Pregnancy or currently planning pregnancy
- Breastfeeding
- Significant comorbidity prohibits use of systemic agents (e.g., liver or kidney disease, blood dyscrasias, uncontrolled hypertension)
- Hypersensitivity
- History of intolerance or adverse event

Appendix B: Risk Factors for Articular Juvenile Idiopathic Arthritis

- Positive rheumatoid factor
- Positive anti-cyclic citrullinated peptide antibodies
- Pre-existing joint damage



AETNA BETTER HEALTH®  
Coverage Policy/Guideline

Name: Enbrel Page: 12 of 14

Effective Date: 2/11/2026 Last Review Date: 1/2026

Applies to:  Illinois  Florida  Florida Kids  
 New Jersey  Maryland  Michigan  
 Pennsylvania Kids  Virginia  Kentucky PRMD

**Approval Duration and Quantity Restrictions:**

**Approval Duration:**

Initial Approval: immune checkpoint inhibitor toxicity when the member has Stevens-Johnson syndrome or toxic epidermal necrolysis = 1 month; all others = 12 months

Renewal Approval: 12 months

**Quantity Level Limit:**

Medication	Standard Limit	Exception Limit*
Enbrel (etanercept) 25 mg per 0.5 mL prefilled syringe	8 syringes per 28 days	None
Enbrel (etanercept) 25 mg /0.5 mL single-dose vial	8 vials per 28 days	16 vials per 28 days
Enbrel (etanercept) 50 mg per 1 mL prefilled syringe/cartridge	4 syringes per 28 days	8 syringes per 28 days
Enbrel (etanercept) 50 mg per mL SureClick Autoinjector	4 cartridges per 28 days	8 cartridges per 28 days

\*Exception limits apply to loading doses.

**References:**

1. Enbrel [package insert]. Thousand Oaks, CA: Immunex Corporation; October 2024.
2. van der Heijde D, Ramiro S, Landewe R, et al. 2016 Update of the international ASAS-EULAR management recommendations for axial spondyloarthritis. Ann Rheum Dis. 2017;0:1-14.
3. Flagg SD, Meador R, Hsia E, et al. Decreased pain and synovial inflammation after etanercept therapy in patients with reactive and undifferentiated arthritis: an open-label trial. Arthritis Rheum. 2005;53(4):613-617.
4. IBM Micromedex® DRUGDEX® System (electronic version). IBM Watson Health, Greenwood Village, Colorado, USA. Available at: <https://www.micromedexsolutions.com/> (cited: 08/14/2025).
5. Smolen JS, Landewé RBM, Bergstra SA, et al. EULAR recommendations for the management of rheumatoid arthritis with synthetic and biological disease-modifying antirheumatic drugs: 2022 update. Ann Rheum Dis. 2023;82:3-18.
6. Singh JA, Saag KG, Bridges SL Jr, et al. 2015 American College of Rheumatology Guideline for the Treatment of Rheumatoid Arthritis. Arthritis Rheumatol. 2016;68(1)1-26.



AETNA BETTER HEALTH®  
Coverage Policy/Guideline

Name: Enbrel Page: 13 of 14

Effective Date: 2/11/2026 Last Review Date: 1/2026

Applies to:  Illinois  Florida  Florida Kids  
 New Jersey  Maryland  Michigan  
 Pennsylvania Kids  Virginia  Kentucky PRMD

7. Saag KG, Teng GG, Patkar NM, et al. American College of Rheumatology 2008 recommendations for the use of nonbiologic and biologic disease-modifying antirheumatic drugs in rheumatoid arthritis. *Arthritis Rheum.* 2008;59(6):762-784.
8. Ringold S, Angeles-Han ST, Beukelman T, et al. 2019 American College of Rheumatology/Arthritis Foundation Guideline for the Treatment of Juvenile Idiopathic Arthritis: Therapeutic Approaches for Non-Systemic Polyarthritis, Sacroiliitis, and Enthesitis. *Arthritis Care Res.* 2019;71(6):717-734. doi:10.1002/acr.23870.
9. Menter A, Gottlieb A, Feldman SR, et al. Guidelines for the management of psoriasis and psoriatic arthritis. Section 1: Overview of psoriasis and guidelines of care for the treatment of psoriasis with biologics. *J Am Acad Dermatol.* 2008;58(5):826-850.
10. Menter A, Korman NJ, Elmets CA, et al. Guidelines of care for the management of psoriasis and psoriatic arthritis. Section 6: Guidelines of care for the treatment of psoriasis and psoriatic arthritis: case-based presentations and evidence-based conclusions. *J Am Acad Dermatol.* 2011;65(1):137-174.
11. Gossec L, Smolen JS, Ramiro S, et al. European League Against Rheumatism (EULAR) recommendations for the management of psoriatic arthritis with pharmacological therapies; 2015 update. *Ann Rheum Dis.* 2016;75(3):499-510.
12. Gladman DD, Antoni C, Mease P, et al. Psoriatic arthritis: epidemiology, clinical features, course, and outcome. *Ann Rheum Dis.* 2005;64(Suppl II):ii14-ii17.
13. Coates LC, Soriano ER, Corp N, et al. Group for Research and Assessment of Psoriasis and Psoriatic Arthritis (GRAPPA): updated treatment recommendations for psoriatic arthritis 2021. *Nat Rev Rheumatol.* 2022;18(8):465-479.
14. Braun J, van den Berg R, Baraliakos X, et al. 2010 update of the ASAS/EULAR recommendations for the management of ankylosing spondylitis. *Ann Rheum Dis.* 2011;70:896-904.
15. Ward MM, Deodhar A, Gensler LS, et al. 2019 Update of the American College of Rheumatology/Spondylitis Association of America/Spondyloarthritis Research and Treatment Network Recommendations for the Treatment of Ankylosing Spondylitis and Nonradiographic Axial Spondyloarthritis. *Arthritis Rheumatol.* 2019;71(10):1599-1613. doi:10.1002/art.41042.
16. Martin PJ, Rizzo JD, Wingard JR, et al. First and second line systemic treatment of acute graft versus host disease: Recommendations of the American Society of Blood and Marrow Transplantation. *Biol Blood Marrow Transplant.* 2012;18(8):1150-1163.
17. Hatemi G, Christensen R, Bang D, et al. 2018 update of the EULAR recommendations for the management of Behcet's syndrome. *Ann Rheum Dis.* 2018;77:808-818.
18. Menter A, Strober BE, Kaplan DH, et al. Joint AAD-NPF guidelines of care for the management and treatment of psoriasis with biologics. *J Am Acad Dermatol.* 2019;80(4):1029-1072.
19. Testing for TB Infection. Centers for Disease Control and Prevention. Retrieved on August 14, 2025 from: <https://www.cdc.gov/tb/testing/index.html>.



AETNA BETTER HEALTH®  
Coverage Policy/Guideline

Name: Enbrel Page: 14 of 14

Effective Date: 2/11/2026 Last Review Date: 1/2026

Applies to:  Illinois  Florida  Florida Kids  
 New Jersey  Maryland  Michigan  
 Pennsylvania Kids  Virginia  Kentucky PRMD

20. Singh JA, Guyatt G, Ogdie A, et al. 2018 American College of Rheumatology/National Psoriasis Foundation Guideline for the Treatment of Psoriatic Arthritis. *Arthritis Rheumatol.* 2019;71(1):5-32. doi:10.1002/art.40726.
21. Flores D, Marquez J, Garza M, Espinoza LR. Reactive arthritis: newer developments. *Rheum Dis Clin North Am.* 2003;29(1):37-vi.
22. The NCCN Drugs & Biologics Compendium 2025 National Comprehensive Cancer Network, Inc. Available at: <http://www.nccn.org>. Accessed August 14, 2025.
23. Lexicomp [database online]. Hudson, OH: Lexi-Comp, Inc.; <https://online.lexi.com/lco/action/home> [available with subscription]. Accessed August 14, 2025.
24. Menter, A, Cordero, KM, Davis, DM, et al. Joint AAD-NPF guidelines of care for the management and treatment of psoriasis in pediatric patients. *J Am Acad Dermatol.* 2020;82(1):161-201.
25. Menter, A, Gelfand, JM, Connor, C, et al. Joint AAD-NPF guidelines of care for the management of psoriasis with systemic nonbiologic therapies. *J Am Acad Dermatol.* 2020;82(6):1445-86.
26. Onel KB, Horton DB, Lovell DJ, et al. 2021 American College of Rheumatology guideline for the treatment of juvenile idiopathic arthritis: therapeutic approaches for oligoarthritis, temporomandibular joint arthritis, and systemic juvenile idiopathic arthritis. *Arthritis Rheumatol.* 2022;74(4):553-569.
27. Elmets C, Korman N, et al. Joint AAD-NPF guidelines of care for the management and treatment of psoriasis with topical therapy and alternative medicine modalities for psoriasis severity measures. *J Am Acad Dermatol.* 2021;84(2):432-470.
28. Alikhan A, Sayed C, Alavi A, et al. North American clinical management guidelines for hidradenitis suppurativa: A publication from the United States and Canadian Hidradenitis Suppurativa Foundations Part I: Diagnosis, evaluation, and the use of complementary and procedural management. *J Am Acad Dermatol.* 2019;81(1):76-90.
29. Alikhan A, Sayed C, Alavi A, et al. North American clinical management guidelines for hidradenitis suppurativa: A publication from the United States and Canadian Hidradenitis Suppurativa Foundations Part II: Topical, intralesional, and systemic medical management. *J Am Acad Dermatol.* 2019;81(1):91-101.
30. Smolen JS, Aletaha D. Assessment of rheumatoid arthritis activity in clinical trials and clinical practice. In: UpToDate, Post TW (Ed), UpToDate, Waltham, MA. Available with subscription. URL: [www.uptodate.com](http://www.uptodate.com). Accessed March 19, 2021.
31. Fraenkel L, Bathon JM, England BR, et al. 2021 American College of Rheumatology guideline for the treatment of rheumatoid arthritis. *Arthrit Care Res.* 2021;0:1-16.
32. Azulfidine [package insert]. New York, NY: Pfizer; October 2022.