



AETNA BETTER HEALTH®
Coverage Policy/Guideline

Name: Humira and Biosimilars Page: 1 of 23

Effective Date: 11/15/2023 Last Review Date: 10/2023

Applies to:	<input checked="" type="checkbox"/> Illinois	<input type="checkbox"/> Florida	<input checked="" type="checkbox"/> Florida Kids
	<input type="checkbox"/> New Jersey	<input type="checkbox"/> Maryland	<input type="checkbox"/> Michigan
	<input checked="" type="checkbox"/> Pennsylvania Kids	<input type="checkbox"/> Virginia	<input checked="" type="checkbox"/> Kentucky PRMD

Intent:

The intent of this policy/guideline is to provide information to the prescribing practitioner outlining the coverage criteria for Humira and biosimilars under the patient's prescription drug benefit.

Description:

A. FDA-Approved Indications

1. Reducing signs and symptoms, inducing major clinical response, inhibiting the progression of structural damage, and improving physical function in adult patients with moderately to severely active rheumatoid arthritis (RA).
2. Reducing signs and symptoms of moderately to severely active polyarticular juvenile idiopathic arthritis (JIA) in patients 2 years of age and older.
3. Reducing signs and symptoms, inhibiting the progression of structural damage, and improving physical function in adult patients with active psoriatic arthritis (PsA).
4. Reducing signs and symptoms in adult patients with active ankylosing spondylitis (AS).
5. The treatment of moderately to severely active Crohn's disease (CD) in adult and pediatric patients 6 years of age and older.
6. The treatment of moderately to severely active ulcerative colitis (UC) in adults and pediatric patients 5 years of age and older.
Limitations of Use: The effectiveness of Humira has not been established in patients who have lost response to or were intolerant to tumor necrosis factor (TNF) blockers.
7. The treatment of adult patients with moderate to severe chronic plaque psoriasis (PsO) who are candidates for systemic therapy or phototherapy, and when other systemic therapies are medically less appropriate.
8. The treatment of moderate to severe hidradenitis suppurativa in patients 12 years of age and older.
9. The treatment of non-infectious intermediate, posterior, and panuveitis in adults and pediatric patients 2 years of age and older.

B. Compendial Uses

1. Axial spondyloarthritis
2. Non-radiographic axial spondyloarthritis
3. Behcet's disease
4. Pyoderma gangrenosum
5. Oligoarticular juvenile idiopathic arthritis
6. Immunotherapy-related inflammatory arthritis



AETNA BETTER HEALTH®
Coverage Policy/Guideline

Name: Humira and Biosimilars Page: 2 of 23

Effective Date: 11/15/2023 Last Review Date: 10/2023

Applies to:	<input checked="" type="checkbox"/> Illinois	<input type="checkbox"/> Florida	<input checked="" type="checkbox"/> Florida Kids
	<input type="checkbox"/> New Jersey	<input type="checkbox"/> Maryland	<input type="checkbox"/> Michigan
	<input checked="" type="checkbox"/> Pennsylvania Kids	<input type="checkbox"/> Virginia	<input checked="" type="checkbox"/> Kentucky PRMD

All other indications are considered experimental/investigational and not medically necessary.

Applicable Drug List:

Preferred:

Humira (adalimumab)

Non-Preferred:

Abrilada (adalimumab-afzb),
Amjevita (adalimumab-atto) Auto-Injector
Amjevita (adalimumab-atto) Prefilled Syringe
Cyltezo (adalimumab-adbm)
Hadlima (adalimumab-bwwd)
Hulio (adalimumab-fkjp)
Hyrimoz (adalimumab-adaz)
Idacio (adalimumab-aacf)
Yuflyma (adalimumab-aaty)
Yusimry (adalimumab-aqvh)
adalimumab-adaz
adalimumab-fkjp

Policy/Guideline:

Documentation for all indications:

For requests for Abrilada (adalimumab-afzb), Amjevita (adalimumab-atto) Auto-Injector, Amjevita (adalimumab-atto) Prefilled Syringe, Cyltezo (adalimumab-adbm), Hadlima (adalimumab-bwwd), Hulio (adalimumab-fkjp), Hyrimoz (adalimumab-adaz), Idacio (adalimumab-aacf), Yuflyma (adalimumab-aaty), Yusimry (adalimumab-aqvh), adalimumab-adaz, or adalimumab-fkjp: the patient is unable to take the preferred adalimumab product, where indicated, for the given diagnosis due to a trial and inadequate treatment response or intolerance, or a contraindication. Documentation is required for approval.

Documentation:

Submission of the following information is necessary to initiate the prior authorization review:

- A. Rheumatoid arthritis (RA)
 - 1. Initial requests:



AETNA BETTER HEALTH®
Coverage Policy/Guideline

Name: Humira and Biosimilars Page: 3 of 23

Effective Date: 11/15/2023 Last Review Date: 10/2023

Applies to:	<input checked="" type="checkbox"/> Illinois	<input type="checkbox"/> Florida	<input checked="" type="checkbox"/> Florida Kids
	<input type="checkbox"/> New Jersey	<input type="checkbox"/> Maryland	<input type="checkbox"/> Michigan
	<input checked="" type="checkbox"/> Pennsylvania Kids	<input type="checkbox"/> Virginia	<input checked="" type="checkbox"/> Kentucky PRMD

- i. Chart notes, medical record documentation, or claims history supporting previous medications tried (if applicable), including response to therapy. If therapy is not advisable, documentation of clinical reason to avoid therapy.
 - ii. Laboratory results, chart notes, or medical record documentation of biomarker testing (i.e., rheumatoid factor [RF], anti-cyclic citrullinated peptide [anti-CCP], and C-reactive protein [CRP] and/or erythrocyte sedimentation rate [ESR]) (if applicable).
 2. Continuation requests: Chart notes or medical record documentation supporting positive clinical response.
- B. Articular juvenile idiopathic arthritis (JIA)
 1. Initial requests: Chart notes, medical record documentation, or claims history supporting previous medications tried (if applicable), including response to therapy.
 2. Continuation requests: Chart notes or medical record documentation supporting positive clinical response.
- C. Ankylosing spondylitis (AS), non-radiographic axial spondyloarthritis (nr-axSpA), psoriatic arthritis (PsA), hidradenitis suppurativa, and uveitis (non-infectious intermediate, posterior and panuveitis)
 1. Initial requests: Chart notes, medical record documentation, or claims history supporting previous medications tried (if applicable), including response to therapy. If therapy is not advisable, documentation of clinical reason to avoid therapy.
 2. Continuation requests: Chart notes or medical record documentation supporting positive clinical response.
- D. Crohn's disease (CD)

Continuation requests: Chart notes or medical record documentation supporting positive clinical response to therapy or remission.
- E. Ulcerative colitis (UC)

Continuation requests: Chart notes or medical record documentation supporting positive clinical response to therapy or remission.
- F. Plaque psoriasis (PsO)
 1. Initial requests:
 - i. Chart notes or medical record documentation of affected area(s) and body surface area (BSA) affected (if applicable).
 - ii. Chart notes, medical record documentation, or claims history supporting previous medications tried (if applicable), including response to therapy. If therapy is not advisable, documentation of clinical reason to avoid therapy.



AETNA BETTER HEALTH®
Coverage Policy/Guideline

Name: Humira and Biosimilars Page: 4 of 23

Effective Date: 11/15/2023 Last Review Date: 10/2023

Applies to:	<input checked="" type="checkbox"/> Illinois	<input type="checkbox"/> Florida	<input checked="" type="checkbox"/> Florida Kids
	<input type="checkbox"/> New Jersey	<input type="checkbox"/> Maryland	<input type="checkbox"/> Michigan
	<input checked="" type="checkbox"/> Pennsylvania Kids	<input type="checkbox"/> Virginia	<input checked="" type="checkbox"/> Kentucky PRMD

2. Continuation requests: Chart notes or medical record documentation of decreased body surface area (BSA) affected and/or improvement in signs and symptoms.

G. Behcet's disease and immunotherapy-related inflammatory arthritis (initial requests only): Chart notes, medical record documentation, or claims history supporting previous medications tried, including response to therapy (if applicable).

H. Pyoderma gangrenosum (initial requests only): Chart notes, medical record documentation, or claims history supporting previous medications tried (if applicable), including response to therapy. If therapy is not advisable, documentation of clinical reason to avoid therapy.

Prescriber Specialty:

This medication must be prescribed by or in consultation with one of the following:

- A. Rheumatoid arthritis, articular juvenile idiopathic arthritis, ankylosing spondylitis, non-radiographic axial spondyloarthritis, and Behcet's disease: rheumatologist
- B. Psoriatic arthritis and hidradenitis suppurativa: rheumatologist or dermatologist
- C. Crohn's disease and ulcerative colitis: gastroenterologist
- D. Plaque psoriasis and pyoderma gangrenosum: dermatologist
- E. Uveitis: ophthalmologist or rheumatologist
- F. Immunotherapy-related inflammatory arthritis: oncologist, hematologist, or rheumatologist

Criteria for Initial Approval:

A. Rheumatoid arthritis (RA)

- 1. Authorization of 12 months may be granted for adult members who have previously received a biologic or targeted synthetic drug (e.g., Rinvoq, Xeljanz) indicated for moderately to severely active rheumatoid arthritis.
- 2. Authorization of 12 months may be granted for adult members for treatment of moderately to severely active RA when all of the following criteria are met:
 - i. Member meets either of the following criteria:
 - a. Member has been tested for either of the following biomarkers and the test was positive:
 - 1. Rheumatoid factor (RF)
 - 2. Anti-cyclic citrullinated peptide (anti-CCP)
 - b. Member has been tested for ALL of the following biomarkers:
 - 1. RF
 - 2. Anti-CCP



AETNA BETTER HEALTH®
Coverage Policy/Guideline

Name: Humira and Biosimilars Page: 5 of 23

Effective Date: 11/15/2023 Last Review Date: 10/2023

Applies to:	<input checked="" type="checkbox"/> Illinois	<input type="checkbox"/> Florida	<input checked="" type="checkbox"/> Florida Kids
	<input type="checkbox"/> New Jersey	<input type="checkbox"/> Maryland	<input type="checkbox"/> Michigan
	<input checked="" type="checkbox"/> Pennsylvania Kids	<input type="checkbox"/> Virginia	<input checked="" type="checkbox"/> Kentucky PRMD

3. C-reactive protein (CRP) and/or erythrocyte sedimentation rate (ESR)
- ii. Member meets either of the following criteria:
 - a. Member has experienced an inadequate response to at least a 3-month trial of methotrexate despite adequate dosing (i.e., titrated to at least 15 mg/week).
 - b. Member has an intolerance or contraindication to methotrexate (see Appendix A).

B. Articular juvenile idiopathic arthritis (JIA)

1. Authorization of 12 months may be granted for members 2 years of age or older who have previously received a biologic or targeted synthetic drug (e.g., Xeljanz) indicated for moderately to severely active articular juvenile idiopathic arthritis.
2. Authorization of 12 months may be granted for members 2 years of age or older for treatment of moderately to severely active articular juvenile idiopathic arthritis when any of the following criteria is met:
 - i. Member has had an inadequate response to methotrexate or another conventional synthetic drug (e.g., leflunomide, sulfasalazine, hydroxychloroquine) administered at an adequate dose and duration.
 - ii. Member has had an inadequate response to a trial of scheduled non-steroidal anti-inflammatory drugs (NSAIDs) and/or intra-articular glucocorticoids (e.g., triamcinolone hexacetonide) and one of the following risk factors for poor outcome:
 - a. Involvement of ankle, wrist, hip, sacroiliac joint, and/or temporomandibular joint (TMJ)
 - b. Presence of erosive disease or enthesitis
 - c. Delay in diagnosis
 - d. Elevated levels of inflammation markers
 - e. Symmetric disease
 - iii. Member has risk factors for disease severity and potentially a more refractory disease course (see Appendix B) and the member also meets one of the following:
 - a. High-risk joints are involved (e.g., cervical spine, wrist, or hip).
 - b. High disease activity.
 - c. Is judged to be at high risk for disabling joint disease.

C. Psoriatic arthritis (PsA)

1. Authorization of 12 months may be granted for adult members who have previously received a biologic or targeted synthetic drug (e.g., Rinvoq, Otezla) indicated for active psoriatic arthritis.



AETNA BETTER HEALTH®
Coverage Policy/Guideline

Name: Humira and Biosimilars Page: 6 of 23

Effective Date: 11/15/2023 Last Review Date: 10/2023

Applies to:	<input checked="" type="checkbox"/> Illinois	<input type="checkbox"/> Florida	<input checked="" type="checkbox"/> Florida Kids
	<input type="checkbox"/> New Jersey	<input type="checkbox"/> Maryland	<input type="checkbox"/> Michigan
	<input checked="" type="checkbox"/> Pennsylvania Kids	<input type="checkbox"/> Virginia	<input checked="" type="checkbox"/> Kentucky PRMD

2. Authorization of 12 months may be granted for adult members for treatment of active psoriatic arthritis when either of the following criteria is met:
 - i. Member has mild to moderate disease and meets one of the following criteria:
 - a. Member has had an inadequate response to methotrexate, leflunomide, or another conventional synthetic drug (e.g., sulfasalazine) administered at an adequate dose and duration.
 - b. Member has an intolerance or contraindication to methotrexate or leflunomide (see Appendix A), or another conventional synthetic drug (e.g., sulfasalazine).
 - c. Member has enthesitis or predominantly axial disease.
 - ii. Member has severe disease.

D. Ankylosing spondylitis (AS) and non-radiographic axial spondyloarthritis (nr-axSpA)

1. Authorization of 12 months may be granted for adult members who have previously received a biologic or targeted synthetic drug (e.g., Rinvoq, Xeljanz) indicated for active ankylosing spondylitis or active non-radiographic axial spondyloarthritis.
2. Authorization of 12 months may be granted for adult members for treatment of active ankylosing spondylitis or active non-radiographic axial spondyloarthritis when either of the following criteria is met:
 - i. Member has experienced an inadequate response to at least two non-steroidal anti-inflammatory drugs (NSAIDs).
 - ii. Member has an intolerance or contraindication to two or more NSAIDs.

E. Crohn's disease (CD)

Authorization of 12 months may be granted for members 6 years of age or older for treatment of moderately to severely active CD.

F. Ulcerative colitis (UC)

Authorization of 12 months may be granted for members 5 years of age or older for treatment of moderately to severely active ulcerative colitis.

G. Plaque psoriasis (PsO)

1. Authorization of 12 months may be granted for adult members who have previously received a biologic or targeted synthetic drug (e.g., Sotyktu, Otezla) indicated for treatment of moderate to severe plaque psoriasis.



AETNA BETTER HEALTH®
Coverage Policy/Guideline

Name: Humira and Biosimilars Page: 7 of 23

Effective Date: 11/15/2023 Last Review Date: 10/2023

Applies to:	<input checked="" type="checkbox"/> Illinois	<input type="checkbox"/> Florida	<input checked="" type="checkbox"/> Florida Kids
	<input type="checkbox"/> New Jersey	<input type="checkbox"/> Maryland	<input type="checkbox"/> Michigan
	<input checked="" type="checkbox"/> Pennsylvania Kids	<input type="checkbox"/> Virginia	<input checked="" type="checkbox"/> Kentucky PRMD

2. Authorization of 12 months may be granted for adult members for treatment of moderate to severe plaque psoriasis when any of the following criteria is met:
 - i. Crucial body areas (e.g., hands, feet, face, neck, scalp, genitals/groin, intertriginous areas) are affected.
 - ii. At least 10% of body surface area (BSA) is affected.
 - iii. At least 3% of body surface area (BSA) is affected and the member meets either of the following criteria:
 - a. Member has had an inadequate response or intolerance to either phototherapy (e.g., UVB, PUVA) or pharmacologic treatment with methotrexate, cyclosporine, or acitretin.
 - b. Member has a clinical reason to avoid pharmacologic treatment with methotrexate, cyclosporine, and acitretin (see Appendix A).

H. Hidradenitis suppurativa

1. Authorization of 12 months may be granted for members 12 years of age or older who have previously received a biologic indicated for treatment of moderate to severe hidradenitis suppurativa.
2. Authorization of 12 months may be granted for member 12 years of age or older for treatment of moderate to severe hidradenitis suppurativa when either of the following is met:
 - i. Member has experienced an inadequate response to an oral antibiotic for at least 90 days.
 - ii. Member has an intolerance or contraindication to oral antibiotics.

I. Uveitis (non-infectious intermediate, posterior and panuveitis)

1. Authorization of 12 months may be granted for members 2 years of age or older who have previously received a biologic indicated for non-infectious intermediate, posterior, and panuveitis.
2. Authorization of 12 months may be granted for members 2 years of age or older for treatment of non-infectious intermediate, posterior and panuveitis when either of the following is met:
 - i. Member has experienced an inadequate response to corticosteroids or immunosuppressive therapy (e.g., azathioprine, cyclosporine, methotrexate).
 - ii. Member has an intolerance or contraindication to corticosteroids and immunosuppressive therapy (e.g., azathioprine, cyclosporine, methotrexate).

J. Behcet's disease



AETNA BETTER HEALTH®
Coverage Policy/Guideline

Name: Humira and Biosimilars Page: 8 of 23

Effective Date: 11/15/2023 Last Review Date: 10/2023

Applies to:	<input checked="" type="checkbox"/> Illinois	<input type="checkbox"/> Florida	<input checked="" type="checkbox"/> Florida Kids
	<input type="checkbox"/> New Jersey	<input type="checkbox"/> Maryland	<input type="checkbox"/> Michigan
	<input checked="" type="checkbox"/> Pennsylvania Kids	<input type="checkbox"/> Virginia	<input checked="" type="checkbox"/> Kentucky PRMD

1. Authorization of 12 months may be granted for members who have previously received Otezla or a biologic indicated for the treatment of Behcet's disease.
2. Authorization of 12 months may be granted for the treatment of Behcet's disease when the member has had an inadequate response to at least one non-biologic medication for Behcet's disease (e.g., apremilast, colchicine, systemic glucocorticoids, azathioprine).

K. Pyoderma gangrenosum

1. Authorization of 12 months may be granted for members who have previously received a biologic indicated for treatment of pyoderma gangrenosum.
2. Authorization of 12 months may be granted for treatment of pyoderma gangrenosum when either of the following is met:
 - i. Member has experienced an inadequate response to corticosteroids or immunosuppressive therapy (e.g., cyclosporine or mycophenolate mofetil).
 - ii. Member has an intolerance or contraindication to corticosteroids and immunosuppressive therapy (e.g., cyclosporine, mycophenolate mofetil).

L. Immunotherapy-related inflammatory arthritis

Authorization of 12 months may be granted for treatment of severe/refractory immunotherapy-related inflammatory arthritis that is not responding to corticosteroids and anti-inflammatory agents.

Continuation of Therapy:

A. Rheumatoid arthritis (RA)

Authorization of 12 months may be granted for all adult members (including new members) who are using the requested medication for moderately to severely active rheumatoid arthritis and who achieve or maintain a positive clinical response as evidenced by disease activity improvement of at least 20% from baseline in tender joint count, swollen joint count, pain, or disability.

B. Articular juvenile idiopathic arthritis (JIA)

Authorization of 12 months may be granted for all members 2 years of age or older (including new members) who are using the requested medication for moderately to severely active articular juvenile idiopathic arthritis and who achieve or maintain a positive clinical response as evidenced by low disease activity or improvement in signs and symptoms of the condition when there is improvement in any of the following from baseline:

1. Number of joints with active arthritis (e.g., swelling, pain, limitation of motion)



AETNA BETTER HEALTH®
Coverage Policy/Guideline

Name: Humira and Biosimilars Page: 9 of 23

Effective Date: 11/15/2023 Last Review Date: 10/2023

Applies to:	<input checked="" type="checkbox"/> Illinois	<input type="checkbox"/> Florida	<input checked="" type="checkbox"/> Florida Kids
	<input type="checkbox"/> New Jersey	<input type="checkbox"/> Maryland	<input type="checkbox"/> Michigan
	<input checked="" type="checkbox"/> Pennsylvania Kids	<input type="checkbox"/> Virginia	<input checked="" type="checkbox"/> Kentucky PRMD

2. Number of joints with limitation of movement
3. Functional ability

C. Psoriatic arthritis (PsA)

Authorization of 12 months may be granted for all adult members (including new members) who are using the requested medication for psoriatic arthritis and who achieve or maintain a positive clinical response as evidenced by low disease activity or improvement in signs and symptoms of the condition when there is improvement in any of the following from baseline:

1. Number of swollen joints
2. Number of tender joints
3. Dactylitis
4. Enthesitis
5. Axial disease
6. Skin and/or nail involvement

D. Ankylosing spondylitis (AS) and non-radiographic axial spondyloarthritis (nr-axSpA)

Authorization of 12 months may be granted for all adult members (including new members) who are using the requested medication for active ankylosing spondylitis or active non-radiographic axial spondyloarthritis and who achieve or maintain a positive clinical response as evidenced by low disease activity or improvement in signs and symptoms of the condition when there is improvement in any of the following from baseline:

1. Functional status
2. Total spinal pain
3. Inflammation (e.g., morning stiffness)

E. Crohn's disease (CD)

1. Authorization of 12 months may be granted for all members 6 years of age or older (including new members) who are using the requested medication for moderately to severely active Crohn's disease and who achieve or maintain remission.
2. Authorization of 12 months may be granted for all members 6 years of age or older (including new members) who are using the requested medication for moderately to severely active Crohn's disease and who achieve or maintain a positive clinical response as evidenced by low disease activity or improvement in signs and symptoms of the condition when there is improvement in any of the following from baseline:
 - i. Abdominal pain or tenderness
 - ii. Diarrhea



AETNA BETTER HEALTH®
Coverage Policy/Guideline

Name: Humira and Biosimilars Page: 10 of 23

Effective Date: 11/15/2023 Last Review Date: 10/2023

Applies to:	<input checked="" type="checkbox"/> Illinois	<input type="checkbox"/> Florida	<input checked="" type="checkbox"/> Florida Kids
	<input type="checkbox"/> New Jersey	<input type="checkbox"/> Maryland	<input type="checkbox"/> Michigan
	<input checked="" type="checkbox"/> Pennsylvania Kids	<input type="checkbox"/> Virginia	<input checked="" type="checkbox"/> Kentucky PRMD

- iii. Body weight
- iv. Abdominal mass
- v. Hematocrit
- vi. Appearance of the mucosa on endoscopy, computed tomography enterography (CTE), magnetic resonance enterography (MRE), or intestinal ultrasound
- vii. Improvement on a disease activity scoring tool (e.g., Crohn's Disease Activity Index [CDAI] score)

F. Ulcerative colitis (UC)

1. Authorization of 12 months may be granted for all members 5 years of age and older (including new members) who are using the requested medication for moderately to severely active ulcerative colitis and who achieve or maintain remission.
2. Authorization of 12 months may be granted for all members 5 years of age and older (including new members) who are using the requested medication for moderately to severely active ulcerative colitis and who achieve or maintain a positive clinical response as evidenced by low disease activity or improvement in signs and symptoms of the condition when there is improvement in any of the following from baseline:
 - i. Stool frequency
 - ii. Rectal bleeding
 - iii. Urgency of defecation
 - iv. C-reactive protein (CRP)
 - v. Fecal calprotectin (FC)
 - vi. Appearance of the mucosa on endoscopy, computed tomography enterography (CTE), magnetic resonance enterography (MRE), or intestinal ultrasound
 - vii. Improvement on a disease activity scoring tool (e.g., Ulcerative Colitis Endoscopic Index of Severity [UCEIS], Mayo score)

G. Plaque psoriasis (PsO)

Authorization of 12 months may be granted for all adult members (including new members) who are using the requested medication for moderate to severe plaque psoriasis and who achieve or maintain a positive clinical response as evidenced by low disease activity or improvement in signs and symptoms of the condition when either of the following is met:

1. Reduction in body surface area (BSA) affected from baseline
2. Improvement in signs and symptoms from baseline (e.g., itching, redness, flaking, scaling, burning, cracking, pain)

H. Hidradenitis suppurativa



AETNA BETTER HEALTH®
Coverage Policy/Guideline

Name: Humira and Biosimilars Page: 11 of 23

Effective Date: 11/15/2023 Last Review Date: 10/2023

Applies to:	<input checked="" type="checkbox"/> Illinois	<input type="checkbox"/> Florida	<input checked="" type="checkbox"/> Florida Kids
	<input type="checkbox"/> New Jersey	<input type="checkbox"/> Maryland	<input type="checkbox"/> Michigan
	<input checked="" type="checkbox"/> Pennsylvania Kids	<input type="checkbox"/> Virginia	<input checked="" type="checkbox"/> Kentucky PRMD

Authorization of 12 months may be granted for all members 12 years of age and older (including new members) who are using the requested medication for moderate to severe hidradenitis suppurativa and who achieve or maintain a positive clinical response as evidenced by low disease activity or improvement in signs and symptoms of the condition when any of the following is met:

1. Reduction in abscess and inflammatory nodule count from baseline
2. Reduced formation of new sinus tracts and scarring
3. Decrease in frequency of inflammatory lesions from baseline
4. Reduction in pain from baseline
5. Reduction in suppuration from baseline
6. Improvement in frequency of relapses from baseline
7. Improvement in quality of life from baseline
8. Improvement on a disease severity assessment tool from baseline

I. Uveitis (non-infectious intermediate, posterior and panuveitis)

Authorization of 12 months may be granted for all members 2 years of age and older (including new members) who are using the requested medication for non-infectious intermediate, posterior, and panuveitis and who achieve or maintain a positive clinical response as evidenced by low disease activity or improvement in signs and symptoms of the condition when the patient meets any of the following:

1. Reduced frequency of disease flares compared to baseline
2. Stability or improvement in anterior chamber (AC) cell grade compared to baseline
3. Stability or improvement in vitreous haze (VH) grade compared to baseline
4. Stability or improvement in visual acuity compared to baseline
5. Reduction in glucocorticoid requirements from baseline
6. No new active inflammatory chorioretinal and/or inflammatory retinal vascular lesions relative to baseline

J. Immunotherapy-related inflammatory arthritis

All members (including new members) requesting authorization for continuation of therapy must meet all initial authorization criteria.

K. All other indications

Authorization of 12 months may be granted for all members (including new members) who are using the requested medication for an indication outlined in Section IV and who achieve or maintain a positive clinical response as evidenced by low disease activity or improvement in signs and symptoms of the condition.

Other:



AETNA BETTER HEALTH®
Coverage Policy/Guideline

Name: Humira and Biosimilars Page: 12 of 23

Effective Date: 11/15/2023 Last Review Date: 10/2023

Applies to:	<input checked="" type="checkbox"/> Illinois	<input type="checkbox"/> Florida	<input checked="" type="checkbox"/> Florida Kids
	<input type="checkbox"/> New Jersey	<input type="checkbox"/> Maryland	<input type="checkbox"/> Michigan
	<input checked="" type="checkbox"/> Pennsylvania Kids	<input type="checkbox"/> Virginia	<input checked="" type="checkbox"/> Kentucky PRMD

For all indications: Member has had a documented negative tuberculosis (TB) test (which can include a tuberculosis skin test [PPD], an interferon-release assay [IGRA], or a chest x-ray)* within 6 months of initiating therapy for persons who are naïve to biologic drugs or targeted synthetic drugs associated with an increased risk of TB.

* If the screening testing for TB is positive, there must be further testing to confirm there is no active disease. Do not administer the requested medication to members with active TB infection. If there is latent disease, TB treatment must be started before initiation of the requested medication.

For all indications: Member cannot use the requested medication concomitantly with any other biologic drug or targeted synthetic drug.

Dosage and Administration:

Approvals may be subject to dosing limits in accordance with FDA-approved labeling, accepted compendia, and/or evidence-based practice guidelines. For rheumatoid arthritis, member must initiate treatment with every other week dosing.

Appendices

Appendix A: Examples of Clinical Reasons to Avoid Pharmacologic Treatment with Methotrexate, Cyclosporine, Acitretin, or Leflunomide

1. Clinical diagnosis of alcohol use disorder, alcoholic liver disease, or other chronic liver disease
2. Drug interaction
3. Risk of treatment-related toxicity
4. Pregnancy or currently planning pregnancy
5. Breastfeeding
6. Significant comorbidity prohibits use of systemic agents (e.g., liver or kidney disease, blood dyscrasias, uncontrolled hypertension)
7. Hypersensitivity
8. History of intolerance or adverse event

Appendix B: Risk Factors for Articular Juvenile Idiopathic Arthritis

1. Positive rheumatoid factor
2. Positive anti-cyclic citrullinated peptide antibodies
3. Pre-existing joint damage



AETNA BETTER HEALTH®
Coverage Policy/Guideline

Name: Humira and Biosimilars Page: 13 of 23

Effective Date: 11/15/2023 Last Review Date: 10/2023

Applies to:	<input checked="" type="checkbox"/> Illinois	<input type="checkbox"/> Florida	<input checked="" type="checkbox"/> Florida Kids
	<input type="checkbox"/> New Jersey	<input type="checkbox"/> Maryland	<input type="checkbox"/> Michigan
	<input checked="" type="checkbox"/> Pennsylvania Kids	<input type="checkbox"/> Virginia	<input checked="" type="checkbox"/> Kentucky PRMD

Approval Duration and Quantity Restrictions:

Approval:

Initial and Renewal Approval: 12 months

Quantity Level Limits:

Abbreviations: RA = rheumatoid arthritis; PsA = psoriatic arthritis; AS = ankylosing spondylitis; PJIA = polyarticular juvenile idiopathic arthritis; CD = Crohn’s disease; UC = ulcerative colitis

Note: The standard limit is designed to allow a quantity sufficient for the most common uses of the medication. The recommended dosing parameters for all FDA-approved indications fall within the standard limits. Coverage of an additional quantity may be reviewed on a case-by-case basis upon request.

Medication	Standard Limit	Exception Limit	FDA-recommended dosing
Humira (adalimumab) 10 mg/0.1 mL single-dose prefilled syringe	2 syringes per 28 days	N/A	<p>RA/PsA/AS</p> <ul style="list-style-type: none"> 40 mg every other week For RA, patients not taking concomitant methotrexate: may increase to 40 mg every week or 80 mg every other week if needed <p>PJIA/Pediatric uveitis (2 years and up)</p> <ul style="list-style-type: none"> 10 kg to < 15 kg: 10 mg every other week 15 kg to < 30 kg: 20 mg every other week ≥ 30 kg: 40 mg every other week <p>Pediatric CD (6 years and up)</p>
Humira (adalimumab) 20 mg/0.2 mL single-dose prefilled syringe	4 syringes per 28 days	N/A	
Humira (adalimumab) 40 mg/0.4 mL single-dose prefilled syringe/pen	4 syringes/pens per 28 days	8 syringes/pens per 14 days	
Humira (adalimumab) 40 mg/0.8 mL single-dose prefilled syringe/pen			
Humira (adalimumab) 80 mg/0.8 mL single-dose prefilled pen	2 pens per 28 days	4 pens per 14 days	
Humira (adalimumab) 80 mg/0.8 mL syringe Pediatric Crohn’s Disease Starter Package	N/A	3 syringes per 28 days	



AETNA BETTER HEALTH®
Coverage Policy/Guideline

Name: Humira and Biosimilars Page: 14 of 23

Effective Date: 11/15/2023 Last Review Date: 10/2023

Applies to:

<input checked="" type="checkbox"/> Illinois	<input type="checkbox"/> Florida	<input checked="" type="checkbox"/> Florida Kids
<input type="checkbox"/> New Jersey	<input type="checkbox"/> Maryland	<input type="checkbox"/> Michigan
<input checked="" type="checkbox"/> Pennsylvania Kids	<input type="checkbox"/> Virginia	<input checked="" type="checkbox"/> Kentucky PRMD

Medication	Standard Limit	Exception Limit	FDA-recommended dosing
Humira (adalimumab) 80 mg/0.8 mL and 40 mg/0.4 mL syringe Pediatric Crohn's Disease Starter Package	N/A	2 syringes per 28 days	<ul style="list-style-type: none"> 17 kg to < 40 kg: loading doses of 80 mg on day 1 and 40 mg two weeks later (day 15); maintenance dose starting at week 4 (day 29) of 20 mg every other week ≥ 40 kg: loading doses of 160 mg on day 1 (given in one day or split over two consecutive days) and 80 mg two weeks later (day 15); maintenance dose starting at week 4 (day 29) of 40 mg every other week <p>Pediatric UC (5 years and up)</p> <ul style="list-style-type: none"> 20 kg to < 40 kg: loading doses of 80 mg on day 1, 40 mg one week later (day 8), and 40 mg one week later (day 15); maintenance dose starting at week 4 (day 29) of 40 mg every other week or 20 mg every week ≥ 40 kg: loading doses of 160 mg on day 1 (single dose or split over two consecutive days), 80 mg one week later (day 8), and 80 mg one week later (day 15); maintenance dose starting at week 4 (day 29) of 80 mg every other week or 40 mg every week <p>Adult CD and UC</p>
Humira (adalimumab) 80 mg/0.8 mL pen Pediatric Ulcerative Colitis Starter Package	N/A	4 pens per 28 days	
Humira (adalimumab) 40 mg/0.8 mL pen Crohn's Disease, Ulcerative Colitis, or Hidradenitis Suppurativa Starter Package	N/A	6 pens per 28 days	
Humira (adalimumab) 80 mg/0.8 mL pen Crohn's Disease, Ulcerative Colitis, or Hidradenitis Suppurativa Starter Package	N/A	3 pens per 28 days	
Humira (adalimumab) 40 mg/0.8 mL pen Psoriasis, Uveitis, or Adolescent Hidradenitis Suppurativa Starter Package	N/A	4 pens per 28 days	
Humira (adalimumab) 80 mg/0.8 mL and 40 mg/0.4 mL pen Psoriasis, Uveitis, or Adolescent Hidradenitis Suppurativa Starter Package	N/A	3 pens per 28 days	



AETNA BETTER HEALTH®
Coverage Policy/Guideline

Name: Humira and Biosimilars

Page: 15 of 23

Effective Date: 11/15/2023

Last Review Date: 10/2023

Applies to:	<input checked="" type="checkbox"/> Illinois	<input type="checkbox"/> Florida	<input checked="" type="checkbox"/> Florida Kids
	<input type="checkbox"/> New Jersey	<input type="checkbox"/> Maryland	<input type="checkbox"/> Michigan
	<input checked="" type="checkbox"/> Pennsylvania Kids	<input type="checkbox"/> Virginia	<input checked="" type="checkbox"/> Kentucky PRMD

Medication	Standard Limit	Exception Limit	FDA-recommended dosing
			<ul style="list-style-type: none"> • Loading doses: 160 mg on day 1 (given in one day or split over two consecutive days), followed by 80 mg two weeks later (day 15) • Maintenance dose: two weeks later (day 29), 40 mg every other week <p>Plaque psoriasis/Adult uveitis</p> <ul style="list-style-type: none"> • 80 mg (day 1), followed by 40 mg every other week starting one week after the initial dose of 80 mg (day 8) <p>Adolescent hidradenitis suppurativa (12 years and up)</p> <ul style="list-style-type: none"> • 30 kg to < 60 kg: 80 mg on day 1, 40 mg on day 8 and subsequent doses 40 mg every other week (day 22) • ≥ 60 kg: Follow adult dosing <p>Adult hidradenitis suppurativa</p> <ul style="list-style-type: none"> • Loading doses: 160 mg on day 1 (given in one day or split over two consecutive days), followed by 80 mg two weeks later (day 15) • Maintenance dose: begin 40 mg every week or 80 mg every other week two weeks later (day 29)



AETNA BETTER HEALTH®
Coverage Policy/Guideline

Name: Humira and Biosimilars Page: 16 of 23

Effective Date: 11/15/2023 Last Review Date: 10/2023

Applies to: Illinois Florida Florida Kids
 New Jersey Maryland Michigan
 Pennsylvania Kids Virginia Kentucky PRMD

Medication	Standard Limit	Exception Limit	FDA-recommended dosing
Abrilada (adalimumab-afzb) 10 mg/0.2 mL single-dose prefilled syringe	2 syringes per 28 days	N/A	Refer to Humira (adalimumab)
Abrilada (adalimumab-afzb) 20 mg/0.4 mL single-dose prefilled syringe	4 syringes per 28 days	N/A	
Abrilada (adalimumab-afzb) 40 mg/0.8 mL single-dose prefilled syringe/pen	4 syringes/pens per 28 days	8 syringes/pens per 14 days	
Amjevita (adalimumab-atto) 10 mg/0.2 mL single-dose prefilled syringe	2 syringes per 28 days	N/A	Refer to Humira (adalimumab)
Amjevita (adalimumab-atto) 20 mg/0.4 mL single-dose prefilled syringe	4 syringes per 28 days	N/A	
Amjevita (adalimumab-atto) 40 mg/0.8 mL single-dose prefilled syringe	4 syringes per 28 days	8 syringes per 14 days	
Amjevita (adalimumab-atto) 40 mg/0.8 mL single-dose SureClick autoinjector	4 autoinjectors per 28 days	8 autoinjectors per 14 days	
Cyltezo (adalimumab-adbm) 10 mg/0.2 mL single-dose prefilled syringe	2 syringes per 28 days	N/A	Refer to Humira (adalimumab)
Cyltezo (adalimumab-adbm) 20 mg/0.4 mL single-dose prefilled syringe	4 syringes per 28 days	N/A	
Cyltezo (adalimumab-adbm) 40 mg/0.8 mL single-dose prefilled syringe/pen auto-injector	4 syringes/pen autoinjectors per 28 days	8 syringes/pen autoinjectors per 14 days	
Cyltezo (adalimumab-adbm) 40 mg/0.8 mL pen auto- injector	N/A	4 pen autoinjectors per 28 days	



AETNA BETTER HEALTH®
Coverage Policy/Guideline

Name: Humira and Biosimilars Page: 17 of 23

Effective Date: 11/15/2023 Last Review Date: 10/2023

Applies to: Illinois Florida Florida Kids
 New Jersey Maryland Michigan
 Pennsylvania Kids Virginia Kentucky PRMD

Medication	Standard Limit	Exception Limit	FDA-recommended dosing
Psoriasis or Uveitis Starter Pack			
Cyltezo (adalimumab-adbm) 40 mg/0.8 mL pen auto-injector Crohn's Disease, Ulcerative Colitis, or Hidradenitis Suppurativa Starter Pack	N/A	6 pen autoinjectors per 28 days	
Hadlima (adalimumab-bwwd) 40 mg/0.4 mL single-dose prefilled syringe/PushTouch auto-injector	4 syringes/ autoinjectors per 28 days	8 syringes/ autoinjectors per 14 days	Refer to Humira (adalimumab)
Hadlima (adalimumab-bwwd) 40 mg/0.8 mL single-dose prefilled syringe/PushTouch auto-injector	4 syringes/ autoinjectors per 28 days	8 syringes/ autoinjectors per 14 days	
Hulio (adalimumab-fkjp) 20 mg/0.4 mL single-dose prefilled syringe	4 syringes per 28 days	N/A	Refer to Humira (adalimumab)
Hulio (adalimumab-fkjp) 40 mg/0.8 mL single-dose prefilled syringe/pen auto-injector	4 syringes/pen autoinjectors per 28 days	8 syringes/pen autoinjectors per 14 days	
adalimumab-fkjp 20 mg/0.4 mL single-dose prefilled syringe	4 syringes per 28 days	N/A	
adalimumab-fkjp 40 mg/0.8 mL single-dose Prefilled syringe/pen auto-injector	4 syringes/pen autoinjectors per 28 days	8 syringes/pen autoinjectors per 14 days	



AETNA BETTER HEALTH®
Coverage Policy/Guideline

Name: Humira and Biosimilars Page: 18 of 23

Effective Date: 11/15/2023 Last Review Date: 10/2023

Applies to: Illinois Florida Florida Kids
 New Jersey Maryland Michigan
 Pennsylvania Kids Virginia Kentucky PRMD

Hyrimoz (adalimumab-adaz) 10 mg/0.1 mL single-dose prefilled syringe	2 syringes per 28 days	N/A	Refer to Humira (adalimumab)
Hyrimoz (adalimumab-adaz) 10 mg/0.2 mL single-dose prefilled syringe			
Hyrimoz (adalimumab-adaz) 20 mg/0.2 mL single-dose prefilled syringe	4 syringes per 28 days	N/A	
Hyrimoz (adalimumab-adaz) 20 mg/0.4 mL single-dose prefilled syringe			
Hyrimoz (adalimumab-adaz) 40 mg/0.4 mL single-dose prefilled syringe/ Sensoready pen auto- injector	4 syringes/ pen autoinjectors per 28 days	8 syringes/ pen autoinjectors per 14 days	
Hyrimoz (adalimumab-adaz) 40 mg/0.8 mL single-dose prefilled syringe/ Sensoready pen auto- injector			
Hyrimoz (adalimumab-adaz) 80 mg/0.8 mL single-dose prefilled syringe/ Sensoready pen auto- injector	2 syringes/ pen autoinjectors per 28 days	4 syringes/ pen autoinjectors per 14 days	
Hyrimoz (adalimumab-adaz) 80 mg/0.8 mL Sensoready pen auto-injector Crohn's disease, Ulcerative Colitis or Hidradenitis Suppurativa Starter Package			
Hyrimoz (adalimumab-adaz) 80 mg/0.8 mL and 40 mg/0.4 mL Sensoready pen auto-injector	N/A	4 pen autoinjectors per 28 days	



AETNA BETTER HEALTH®
Coverage Policy/Guideline

Name: Humira and Biosimilars Page: 19 of 23

Effective Date: 11/15/2023 Last Review Date: 10/2023

Applies to: Illinois Florida Florida Kids
 New Jersey Maryland Michigan
 Pennsylvania Kids Virginia Kentucky PRMD

Crohn's disease, Ulcerative Colitis or Hidradenitis Suppurativa Starter Package			
Hyrimoz (adalimumab-adaz) 80 mg/0.8 mL and 40 mg/0.4 mL Sensoready pen auto-injector Plaque Psoriasis Starter Package	N/A	3 pen autoinjectors per 28 days	
Hyrimoz (adalimumab-adaz) 80 mg/0.8 mL prefilled syringe Pediatric Crohn's Disease Starter Pack	N/A	3 syringes per 28 days	Refer to Humira (adalimumab)
Hyrimoz (adalimumab-adaz) 80 mg/0.8 mL and 40 mg/0.4 mL prefilled syringe Pediatric Crohn's Disease Starter Pack	N/A	2 syringes per 28 days	
adalimumab-adaz 40 mg/0.4 mL single-dose prefilled syringe/ Sensoready pen auto-injector	4 syringes/ pen autoinjectors per 28 days	8 syringes/ pen autoinjectors per 14 days	
Idacio (adalimumab-aacf) 40 mg/0.8 mL single-dose prefilled syringe/pen auto-injector	4 syringes/pen autoinjectors per 28 days	8 syringes/pen autoinjectors per 14 days	Refer to Humira (adalimumab)
Idacio (adalimumab-aacf) 40 mg/0.8 mL pen auto-injector Plaque Psoriasis Starter Package	N/A	4 pen autoinjectors (2 trays) per 28 days	
Idacio (adalimumab-aacf) 40 mg/0.8 mL pen auto-injector	N/A	6 pen autoinjectors	



AETNA BETTER HEALTH®
Coverage Policy/Guideline

Name: Humira and Biosimilars Page: 20 of 23

Effective Date: 11/15/2023 Last Review Date: 10/2023

Applies to: Illinois Florida Florida Kids
 New Jersey Maryland Michigan
 Pennsylvania Kids Virginia Kentucky PRMD

Crohn's Disease or Ulcerative Colitis Starter Package		(3 trays) per 28 days	
Yuflyma (adalimumab-aaty) 40 mg/0.4 mL single-dose prefilled syringe/auto-injector	4 syringes/ autoinjectors per 28 days	8 syringes/ autoinjectors per 14 days	Refer to Humira (adalimumab)
Yusimry (adalimumab-aqvh) 40 mg/0.8 mL single-dose prefilled syringe/pen	4 syringes/pens per 28 days	8 syringes/pens per 14 days	Refer to Humira (adalimumab)

References:

- Humira [package insert]. North Chicago, IL: AbbVie Inc.; February 2021.
- Abrilada [package insert]. New York, NY: Pfizer Inc.; June 2023.
- adalimumab-adaz [package insert]. Princeton, NJ: Sandoz Inc.; March 2023.
- adalimumab-fkjp [package insert]. Morgantown, WV: Mylan Specialty L.P.; June 2023.
- Amjevita [package insert]. Thousand Oaks, CA: Amgen Inc., April 2023.
- Cyltezo [package insert]. Ridgefield, CT: Boehringer Ingelheim Pharmaceuticals, Inc.; June 2023.
- Hadlima [package insert]. Jersey City, NJ: Organon & Co.; June 2023.
- Hulio [package insert]. Morgantown, WV: Mylan Specialty L.P.; March 2023.
- Hyrimoz [package insert]. Princeton, NJ: Sandoz Inc.; April 2023.
- Idacio [package insert]. Lake Zurich, IL: Fresenius Kabi USA, LLC, December 2022.
- Yuflyma [package insert]. Jersey City, NJ: Celltrion USA, Inc.; May 2023.
- Yusimry [package insert]. Redwood City, CA: Coherus BioSciences, Inc.; March 2023.
- van der Heijde D, Ramiro S, Landewe R, et al. 2016 Update of the international ASAS-EULAR management recommendations for axial spondyloarthritis. *Ann Rheum Dis.* 2017;0:1-14.
- Sieper J, van der Heijde D, Dougados M, et al. Efficacy and safety of adalimumab in patients with non-radiographic axial spondyloarthritis: results of a randomised placebo-controlled trial (ABILITY-1). *Ann Rheum Dis.* 2013;72(6):815-22.
- Smolen JS, Landewé R, Billsma J, et al. EULAR recommendations for the management of rheumatoid arthritis with synthetic and biological disease-modifying antirheumatic drugs: 2016 update. *Ann Rheum Dis.* 2017;0:1-18.
- Singh JA, Saag KG, Bridges SL Jr, et al. 2015 American College of Rheumatology Guideline for the Treatment of Rheumatoid Arthritis. *Arthritis Rheumatol.* 2016;68(1):1-26.
- Saag KG, Teng GG, Patkar NM, et al. American College of Rheumatology 2008 recommendations for the use of nonbiologic and biologic disease-modifying antirheumatic drugs in rheumatoid arthritis. *Arthritis Rheum.* 2008;59(6):762-784.



AETNA BETTER HEALTH®
Coverage Policy/Guideline

Name: Humira and Biosimilars Page: 21 of 23

Effective Date: 11/15/2023 Last Review Date: 10/2023

Applies to:	<input checked="" type="checkbox"/> Illinois	<input type="checkbox"/> Florida	<input checked="" type="checkbox"/> Florida Kids
	<input type="checkbox"/> New Jersey	<input type="checkbox"/> Maryland	<input type="checkbox"/> Michigan
	<input checked="" type="checkbox"/> Pennsylvania Kids	<input type="checkbox"/> Virginia	<input checked="" type="checkbox"/> Kentucky PRMD

18. Ringold S, Angeles-Han ST, Beukelman T, et al. 2019 American College of Rheumatology/Arthritis Foundation Guideline for the Treatment of Juvenile Idiopathic Arthritis: Therapeutic Approaches for Non-Systemic Polyarthritis, Sacroiliitis, and Enthesitis. *Arthritis Care Res.* 2019;71(6):717-734. doi:10.1002/acr.23870.
19. Gossec L, Baraliakos X, Kerschbaumer A, et al. European League Against Rheumatism (EULAR) recommendations for the management of psoriatic arthritis with pharmacological therapies; 2019 update. *Ann Rheum Dis.* 2020;79(6):700-712.
20. Gladman DD, Antoni C, Mease P, et al. Psoriatic arthritis: epidemiology, clinical features, course, and outcome. *Ann Rheum Dis* 2005;64(Suppl II):ii14-ii17.
21. Coates LC, Soriano ER, Corp N, et al. Group for Research and Assessment of Psoriasis and Psoriatic Arthritis (GRAPPA): updated treatment recommendations for psoriatic arthritis 2021. *Nat Rev Rheumatol.* 2022;18(8):465-479.
22. Menter A, Korman NJ, Elmets CA, et al. Guidelines of care for the management of psoriasis and psoriatic arthritis. Section 6: Guidelines of care for the treatment of psoriasis and psoriatic arthritis: case-based presentations and evidence-based conclusions. *J Am Acad Dermatol.* 2011;65(1):137-174.
23. Braun J, van den Berg R, Baraliakos X, et al. 2010 update of the ASAS/EULAR recommendations for the management of ankylosing spondylitis. *Ann Rheum Dis* 2011;70:896-904.
24. Ward MM, Deodhar A, Gensler LS, et al. 2019 Update of the American College of Rheumatology/Spondylitis Association of America/Spondyloarthritis Research and Treatment Network Recommendations for the Treatment of Ankylosing Spondylitis and Nonradiographic Axial Spondyloarthritis. *Arthritis Rheumatol.* 2019;71(10):1599-1613. doi:10.1002/art.41042.
25. Talley NJ, Abreu MT, Achkar J, et al. An evidence-based systematic review on medical therapies for inflammatory bowel disease. *Am J Gastroenterol.* 2011;106(Suppl 1):S2-S25.
26. Lichtenstein GR, Loftus Jr EV, Isaacs KI, et al. ACG Clinical Guideline: Management of Crohn's Disease in Adults. *Am J Gastroenterol.* 2018;113:481-517.
27. Agarwal A, Andrews JM. Systematic review: IBD-associated pyoderma gangrenosum in the biologic era, the response to therapy. *Aliment Pharmacol Ther.* 2013;38(6):563-572.
28. Arguelles-Arias F, Castro-Laria L, Lobaton T, et al. Characteristics and treatment of pyoderma gangrenosum in inflammatory bowel disease. *Dig Dis Sci.* 2013;58(10):2949-2954.
29. Marzano AV, Ishak RS, Saibeni S, et al. Autoinflammatory skin disorders in inflammatory bowel diseases, pyoderma gangrenosum and Sweet's syndrome: A comprehensive review and disease classification criteria. *Clin Rev Allergy Immunol.* 2013;45(2):202-210.
30. Hatemi G, Christensen R, Bodaghi, et al. 2018 update of the EULAR recommendations for the management of Behcet's syndrome. *Ann Rheum Dis.* 2018.; 77: 808-818.
31. Menter A, Strober BE, Kaplan DH, et al. Joint AAD-NPF guidelines of care for the management and treatment of psoriasis with biologics. *J Am Acad Dermatol.* 2019;80(4):1029-1072.
32. Testing for TB Infection. Centers for Disease Control and Prevention. Retrieved on August 9, 2022 from: <https://www.cdc.gov/tb/topic/testing/tbtesttypes.htm>.
33. Singh JA, Guyatt G, Ogdie A, et al. 2018 American College of Rheumatology/National Psoriasis Foundation Guideline for the Treatment of Psoriatic Arthritis. *Arthritis Rheumatol.* 2019;71(1):5-32. doi:10.1002/art.40726.
34. Feuerstein JD, Isaacs KL, Schneider Y, et al. AGA Clinical Practice Guidelines on the Management of Moderate to Severe Ulcerative Colitis. *Gastroenterology.* 2020; 158:1450.



AETNA BETTER HEALTH®
Coverage Policy/Guideline

Name: Humira and Biosimilars Page: 22 of 23

Effective Date: 11/15/2023 Last Review Date: 10/2023

Applies to:	<input checked="" type="checkbox"/> Illinois	<input type="checkbox"/> Florida	<input checked="" type="checkbox"/> Florida Kids
	<input type="checkbox"/> New Jersey	<input type="checkbox"/> Maryland	<input type="checkbox"/> Michigan
	<input checked="" type="checkbox"/> Pennsylvania Kids	<input type="checkbox"/> Virginia	<input checked="" type="checkbox"/> Kentucky PRMD

35. Menter, A, Cordero, KM, Davis, DM, et al. Joint AAD-NPF guidelines of care for the management and treatment of psoriasis in pediatric patients. *J Am Acad Dermatol.* 2020;82(1):161-201.
36. Menter A, Gelfand JM, Connor C, et al. Joint AAD-NPF guidelines of care for the management of psoriasis with systemic nonbiologic therapies. *J Am Acad Dermatol.* 2020;82(6):1445-86.
37. Micromedex® (electronic version). IBM Watson Health, Greenwood Village, Colorado, USA. Available at: <https://www.micromedexsolutions.com>. Accessed August 20, 2022.
38. George C, Deroide F, Rustin M. Pyoderma gangrenosum – a guide to diagnosis and management. *Clin Med.* 2019;19(3): 224-8.
39. Jaffe G, Dick A, Brezin A, et al. Adalimumab in Patients with Active Noninfectious Uveitis. *N Engl J Med.* 2016; 375: 932-943.
40. Nguyen QD, Merrill P, Jaffe G, et al. Adalimumab for prevention of uveitic flare in patients with inactive non-infectious uveitis controlled by corticosteroids (VISUAL II): a multicentre, double-masked, randomized, placebo-controlled phase 3 trial. *Lancet.* 2016; 388(10050):1183-92.
41. Aletaha D, Neogi T, Silman AJ, et al. 2010 Rheumatoid arthritis classification criteria: an American College of Rheumatology/European League Against Rheumatism collaborative initiative. *Arthritis Rheum.* 2010;62(9):2569-81.
42. Rubin DT, Ananthakrishnan AN, et al. 2019 ACG Clinical Guideline: Ulcerative Colitis in Adults. *Am J Gastroenterol.* 2019;114:384-413.
43. Alikhan A, Sayed C, Alavi A, et al. North American clinical management guidelines for hidradenitis suppurativa: A publication from the United States and Canadian Hidradenitis Suppurativa Foundations Part I: Diagnosis, evaluation, and the use of complementary and procedural management. *J Am Acad Dermatol.* 2019; 81(1): 76-90.
44. Alikhan A, Sayed C, Alavi A, et al. North American clinical management guidelines for hidradenitis suppurativa: A publication from the United States and Canadian Hidradenitis Suppurativa Foundations Part II: Topical, intralesional, and systemic medical management. *J Am Acad Dermatol.* 2019; 81(1): 91-101.
45. Smolen JS, Aletaha D. Assessment of rheumatoid arthritis activity in clinical trials and clinical practice. In: UpToDate, Post TW (Ed), UpToDate, Waltham, MA. Available with subscription. URL: www.uptodate.com. Accessed March 19, 2021.
46. The NCCN Drugs & Biologics Compendium 2022 National Comprehensive Cancer Network, Inc. Available at: <http://www.nccn.org>. Accessed August 15, 2022.
47. Feuerstein J, Ho E, Shmidt E, et al. AGA Clinical Practice Guidelines on the Medical Management of Moderate to Severe Luminal and Perianal Fistulizing Crohn's Disease. *Gastroenterology.* 2021; 160:2496-2508.
48. Fraenkel L, Bathon JM, England BR, et al. 2021 American College of Rheumatology guideline for the treatment of rheumatoid arthritis. *Arthrit Care Res.* 2021;0:1-16.
49. Elmets C, Korman N, et al. Joint AAD-NPF guidelines of care for the management and treatment of psoriasis with topical therapy and alternative medicine modalities for psoriasis severity measures. *J Am Acad Dermatol.* 2021; 84(2):432-470.
50. Onel KB, Horton DB, Lovell DJ, et al. 2021 American College of Rheumatology guideline for the treatment of juvenile idiopathic arthritis: therapeutic approaches for oligoarthritis, temporomandibular joint arthritis, and systemic juvenile idiopathic arthritis. *Arthritis Rheumatol.* 2022;74(4):553-569.



AETNA BETTER HEALTH®
Coverage Policy/Guideline

Name: Humira and Biosimilars Page: 23 of 23

Effective Date: 11/15/2023 Last Review Date: 10/2023

Applies to:	<input checked="" type="checkbox"/> Illinois	<input type="checkbox"/> Florida	<input checked="" type="checkbox"/> Florida Kids
	<input type="checkbox"/> New Jersey	<input type="checkbox"/> Maryland	<input type="checkbox"/> Michigan
	<input checked="" type="checkbox"/> Pennsylvania Kids	<input type="checkbox"/> Virginia	<input checked="" type="checkbox"/> Kentucky PRMD

51. Angeles-Han ST, Ringold S, Beukelman T, et al. 2019 American College of Rheumatology/Arthritis Foundation guideline for the screening, monitoring, and treatment of juvenile idiopathic arthritis-associated uveitis. *Arthritis Care Res.* 2019; 71(6):703-716.