



AETNA BETTER HEALTH®  
Coverage Policy/Guideline

Name: Kineret

Page: 1 of 10

Effective Date: 2/2/2026

Last Review Date: 12/2025

Applies to:	<input type="checkbox"/> Illinois	<input type="checkbox"/> Florida	<input type="checkbox"/> Florida Kids
	<input checked="" type="checkbox"/> New Jersey	<input type="checkbox"/> Maryland	<input type="checkbox"/> Michigan
	<input type="checkbox"/> Pennsylvania Kids	<input type="checkbox"/> Virginia	<input type="checkbox"/> Kentucky PRMD

### Intent:

The intent of this policy/guideline is to provide information to the prescribing practitioner outlining the coverage criteria for Kineret under the patient's prescription drug benefit.

### Description:

The indications below including FDA-approved indications and compendial uses are considered a covered benefit provided that all the approval criteria are met and the member has no exclusions to the prescribed therapy.

#### FDA-Approved Indications<sup>1</sup>

- Moderately to severely active rheumatoid arthritis (RA), in patients 18 years of age or older who have failed 1 or more disease modifying antirheumatic drugs (DMARDs) – (Reference NJ Protocol for Cryopyrin-Associated Periodic Syndromes (CAPS) Products)
- Cryopyrin-Associated Periodic Syndromes (CAPS), including Neonatal-Onset Multisystem Inflammatory Disease (NOMID) – (Reference NJ Protocol for Cryopyrin-Associated Periodic Syndromes (CAPS) Products)
- Deficiency of Interleukin-1 Receptor Antagonist (DIRA)

#### Compendial Uses

- Systemic juvenile idiopathic arthritis (sJIA)<sup>3</sup> – (Reference NJ Protocol for Cryopyrin-Associated Periodic Syndromes (CAPS) Products)
- Adult-onset Still's disease (AOSD)<sup>4-5</sup>
- Multicentric Castleman disease<sup>6</sup>
- Recurrent pericarditis (RP)<sup>16</sup>
- Hyperimmunoglobulin D syndrome (HIDS)/Mevalonate Kinase Deficiency (MKD)<sup>17-19</sup>
- Schnitzler syndrome<sup>2,20,21</sup> – (Reference NJ Protocol for Cryopyrin-Associated Periodic Syndromes (CAPS) Products)
- Gout and pseudogout (calcium pyrophosphate deposition)<sup>2,25-27</sup>
- Chimeric antigen receptor (CAR) T-cell-related toxicities<sup>6</sup>
- Erdheim-Chester Disease<sup>6</sup>
- Immune checkpoint inhibitor-related toxicity—immunotherapy-related hemophagocytic lymphohistiocytosis (HLH)-like syndrome<sup>6</sup>

All other indications are considered experimental/investigational and not medically necessary.



AETNA BETTER HEALTH®  
Coverage Policy/Guideline

Name: Kineret Page: 2 of 10

Effective Date: 2/2/2026 Last Review Date: 12/2025

Applies to:	<input type="checkbox"/> Illinois	<input type="checkbox"/> Florida	<input type="checkbox"/> Florida Kids
	<input checked="" type="checkbox"/> New Jersey	<input type="checkbox"/> Maryland	<input type="checkbox"/> Michigan
	<input type="checkbox"/> Pennsylvania Kids	<input type="checkbox"/> Virginia	<input type="checkbox"/> Kentucky PRMD

**Applicable Drug List:**

Non-preferred:

Kineret

**Policy/Guideline:**

**Documentation for all indications:**

The patient is unable to take TWO preferred products (a preferred adalimumab product, a preferred tocilizumab product, Enbrel, Kevzara or Rinvoq), where indicated, for the given diagnosis due to a trial and inadequate treatment response or intolerance, or a contraindication. Documentation is required for approval.

**Documentation**

Submission of the following information is necessary to initiate the prior authorization review:

Adult-Onset Still's Disease (AOSD)

Initial requests

Chart notes, medical record documentation, or claims history supporting previous medications tried (if applicable).

Continuation requests

Chart notes or medical record documentation supporting positive clinical response.

Neonatal-Onset Multisystem Inflammatory Disease (NOMID)

Continuation requests: Chart notes, medical record documentation, or laboratory results supporting positive clinical response.

Deficiency of Interleukin-1 Receptor Antagonist (DIRA)

Initial requests: IL1RN gene variant status

Recurrent Pericarditis (RP)

Initial requests

Chart notes, medical record documentation, or claims history supporting previous medications tried, including response to therapy.

Continuation requests

Chart notes or medical record documentation supporting positive clinical response.



AETNA BETTER HEALTH®  
Coverage Policy/Guideline

Name: Kineret Page: 3 of 10

Effective Date: 2/2/2026 Last Review Date: 12/2025

Applies to:	<input type="checkbox"/> Illinois	<input type="checkbox"/> Florida	<input type="checkbox"/> Florida Kids
	<input checked="" type="checkbox"/> New Jersey	<input type="checkbox"/> Maryland	<input type="checkbox"/> Michigan
	<input type="checkbox"/> Pennsylvania Kids	<input type="checkbox"/> Virginia	<input type="checkbox"/> Kentucky PRMD

### Hyperimmunoglobulin D Syndrome (HIDS)/Mevalonate Kinase Deficiency (MKD)

Initial requests: Chart notes, medical record documentation, or laboratory result (if applicable) indicating number of active flares within the last 6 months and Physician's Global Assessment (PGA) score or C-reactive protein (CRP) level.

### Gout and Pseudogout Flares, CAR T-Cell-Related Toxicities, and Immune Checkpoint Inhibitor-Related Toxicity

Initial requests: Chart notes, medical record documentation, or claims history supporting previous medications tried (if applicable), including response to therapy. If therapy is not advisable, documentation of clinical reason to avoid therapy.

## **Prescriber Specialties**

This medication must be prescribed by or in consultation with one of the following:

- Adult-onset Still's disease (AOSD), gout, and pseudogout: rheumatologist
- Deficiency of interleukin-1 receptor antagonist (DIRA), and hyperimmunoglobulin D syndrome (HIDS)/mevalonate kinase deficiency (MKD): rheumatologist or immunologist
- Recurrent pericarditis (RP): cardiologist, rheumatologist, or immunologist
- Multicentric Castleman disease, CAR T-cell-related toxicities, Erdheim-Chester disease, and immune checkpoint inhibitor-related toxicity: oncologist or hematologist

## **Coverage Criteria**

### Adult-Onset Still's Disease (AOSD)<sup>4,5,14,15,23,24,34</sup>

Authorization of 12 months may be granted for members who have previously received a biologic indicated for active AOSD.

Authorization of 12 months may be granted for treatment of active AOSD when both of the following criteria are met:

- Member has active systemic features (e.g., fever, arthralgia/arthritis, evanescent rash, lymphadenopathy, hepatomegaly, splenomegaly, sore throat).
- Member meets any of the following:
  - Member has had an inadequate response to a trial of nonsteroidal anti-inflammatory drugs (NSAIDs).
  - Member has had an inadequate response to a trial of corticosteroids.



AETNA BETTER HEALTH®  
Coverage Policy/Guideline

Name: Kineret Page: 4 of 10

Effective Date: 2/2/2026 Last Review Date: 12/2025

Applies to:	<input type="checkbox"/> Illinois	<input type="checkbox"/> Florida	<input type="checkbox"/> Florida Kids
	<input checked="" type="checkbox"/> New Jersey	<input type="checkbox"/> Maryland	<input type="checkbox"/> Michigan
	<input type="checkbox"/> Pennsylvania Kids	<input type="checkbox"/> Virginia	<input type="checkbox"/> Kentucky PRMD

- Member has had an inadequate response to a trial of a conventional synthetic drug (e.g., methotrexate).

#### Neonatal-Onset Multisystem Inflammatory Disease (NOMID)<sup>1</sup>

Authorization of 12 months may be granted for treatment of cryopyrin-associated periodic syndromes (CAPS), including NOMID (also known as chronic infantile neurologic cutaneous and articular [CINCA] syndrome).

#### Deficiency of Interleukin-1 Receptor Antagonist (DIRA)<sup>1,29</sup>

Authorization of 12 months may be granted for treatment of genetically confirmed deficiency of interleukin-1 receptor antagonist (DIRA) due to IL1RN gene variants.

#### Recurrent Pericarditis (RP)<sup>16,33,37</sup>

Authorization of 12 months may be granted for treatment of recurrent pericarditis when both of the following criteria are met:

- Member has had at least two episodes of pericarditis.
- Member has failed at least two agents of standard therapy (e.g., colchicine, non-steroidal anti-inflammatory drugs [NSAIDs], corticosteroids).

#### Multicentric Castleman Disease<sup>6</sup>

Authorization of 12 months may be granted for treatment of multicentric Castleman disease when both of the following criteria are met:

- The requested medication will be used as a single agent.
- The disease has progressed following treatment of relapsed/refractory or progressive disease.

#### Hyperimmunoglobulin D Syndrome (HIDS)/Mevalonate Kinase Deficiency (MKD)<sup>17-19</sup>

Authorization of 12 months may be granted for treatment of HIDS/MKD when both of the following criteria are met:

- Member has had active flares within the last 6 months.
- Physician's Global Assessment (PGA) score greater than or equal to 2 or C-reactive protein (CRP) greater than 10 mg/L.

#### Gout and Pseudogout Flares<sup>25-38</sup>

Authorization of 12 months may be granted for adult members for treatment of flares for gout and pseudogout (also known as calcium pyrophosphate deposition disease) when both of the following criteria are met:

- Member has experienced at least three gout flares in the last 12 months.



AETNA BETTER HEALTH®  
Coverage Policy/Guideline

Name: Kineret Page: 5 of 10

Effective Date: 2/2/2026 Last Review Date: 12/2025

Applies to:	<input type="checkbox"/> Illinois	<input type="checkbox"/> Florida	<input type="checkbox"/> Florida Kids
	<input checked="" type="checkbox"/> New Jersey	<input type="checkbox"/> Maryland	<input type="checkbox"/> Michigan
	<input type="checkbox"/> Pennsylvania Kids	<input type="checkbox"/> Virginia	<input type="checkbox"/> Kentucky PRMD

- Member has had an inadequate response, intolerance, or contraindication to non-steroidal anti-inflammatory drugs (NSAIDs), colchicine, and corticosteroids.

#### Chimeric Antigen Receptor (CAR) T-Cell-Related Toxicities<sup>6</sup>

Authorization of 1 month may be granted for management of chimeric antigen receptor (CAR) T-cell-induced cytokine release syndrome when either of the following criteria is met:

- Cytokine release syndrome is refractory to high-dose corticosteroids and anti-IL-6 therapy.
- Kineret will be used as a replacement for the second dose of tocilizumab when supplies are limited or unavailable.

Authorization of 3 months may be granted for the prophylaxis of CAR T-cell-induced toxicity in members at high risk of developing high-grade immune effector cell-associated neurotoxicity syndrome (ICANS).

#### Erdheim-Chester Disease<sup>6</sup>

Authorization of 12 months may be granted for the treatment of Erdheim-Chester disease.

#### Immune Checkpoint Inhibitor-Related Toxicity<sup>6</sup>

Authorization of 6 months may be granted for the treatment of immune checkpoint inhibitor-related toxicity when the member has immunotherapy-related hemophagocytic lymphohistiocytosis (HLH)-like syndrome and either of the following criteria is met:

- Member has had an inadequate response to systemic corticosteroids.
- Member has an intolerance or contraindication to corticosteroids.

### **Continuation of Therapy**

#### Adult-Onset Still's Disease (AOSD)<sup>3-5,12-15,23,24,34</sup>

Authorization of 12 months may be granted for all members (including new members) who are using the requested medication for adult-onset Still's disease or systemic juvenile idiopathic arthritis and who achieve or maintain a positive clinical response as evidenced by low disease activity or improvement in signs and symptoms of the condition when there is improvement in any of the following from baseline:

- Number of joints with active arthritis (e.g., swelling, pain, limitation of motion)
- Number of joints with limitation of movement
- Functional ability
- Systemic features (e.g., fever, evanescent rash, lymphadenopathy, hepatomegaly, splenomegaly, serositis)



AETNA BETTER HEALTH®  
Coverage Policy/Guideline

Name: Kineret Page: 6 of 10

Effective Date: 2/2/2026 Last Review Date: 12/2025

Applies to:	<input type="checkbox"/> Illinois	<input type="checkbox"/> Florida	<input type="checkbox"/> Florida Kids
	<input checked="" type="checkbox"/> New Jersey	<input type="checkbox"/> Maryland	<input type="checkbox"/> Michigan
	<input type="checkbox"/> Pennsylvania Kids	<input type="checkbox"/> Virginia	<input type="checkbox"/> Kentucky PRMD

### Neonatal-Onset Multisystem Inflammatory Disease (NOMID)<sup>1,30</sup>

Authorization of 12 months may be granted for all members (including new members) who are using the requested medication for CAPS, including NOMID (also known as CINCA), and who achieve or maintain a positive clinical response as evidenced by low disease activity or improvement in signs and symptoms of the condition when there is improvement in any of the following from baseline:

- Fever
- Skin rash
- Joint pain and/or inflammation
- Central nervous system (CNS) symptoms (e.g., meningitis, headache, cerebral atrophy, uveitis, hearing loss)
- Inflammatory markers (e.g., serum amyloid A [SAA], C-reactive protein [CRP], erythrocyte sedimentation rate [ESR])

### Recurrent Pericarditis (RP)<sup>16,33</sup>

Authorization of 12 months may be granted for all members (including new members) who are using the requested medication for recurrent pericarditis and who achieve or maintain a positive clinical response as evidenced by decreased recurrence of pericarditis or improvement in signs and symptoms of the condition when there is improvement in any of the following:

- Pericarditic or pleuritic chest pain
- Pericardial or pleural rubs
- Electrocardiogram (ECG)
- Pericardial effusion
- C-reactive protein (CRP)

### Multicentric Castleman Disease

Authorization of 12 months may be granted for continued treatment of multicentric Castleman disease in members requesting reauthorization who have not experienced disease progression or an unacceptable toxicity.

### CAR T-Cell-Related Toxicities and Immune Checkpoint Inhibitor-Related Toxicity

All members (including new members) requesting authorization for continuation of therapy must meet all requirements in the coverage criteria.

### All Other Indications

Authorization of 12 months may be granted for all members (including new members) who are using the requested medication for an indication outlined in the coverage criteria and



AETNA BETTER HEALTH®  
Coverage Policy/Guideline

Name: Kineret Page: 7 of 10

Effective Date: 2/2/2026 Last Review Date: 12/2025

Applies to:	<input type="checkbox"/> Illinois	<input type="checkbox"/> Florida	<input type="checkbox"/> Florida Kids
	<input checked="" type="checkbox"/> New Jersey	<input type="checkbox"/> Maryland	<input type="checkbox"/> Michigan
	<input type="checkbox"/> Pennsylvania Kids	<input type="checkbox"/> Virginia	<input type="checkbox"/> Kentucky PRMD

who achieve or maintain a positive clinical response as evidenced by low disease activity or improvement in signs and symptoms of the condition.

**Other<sup>1,22</sup>**

For all indications: Member has had a documented negative tuberculosis (TB) test (which can include a tuberculosis skin test [TST] or an interferon-release assay [IGRA]) within 12 months of initiating therapy for persons who are naïve to biologic drugs or targeted synthetic drugs associated with an increased risk of TB.

If the screening testing for TB is positive, there must be further testing to confirm there is no active disease (e.g., chest x-ray). Do not administer the requested medication to members with active TB infection. If there is latent disease, TB treatment must be started before initiation of the requested medication.

For all indications: Member cannot use the requested medication concomitantly with any other biologic drug or targeted synthetic drug.

**Dosage and Administration**

Approvals may be subject to dosing limits in accordance with FDA-approved labeling, accepted compendia, and/or evidence-based practice guidelines.

**Appendix**

Examples of Clinical Reasons to Avoid Pharmacologic Treatment with Methotrexate, Hydroxychloroquine, or Leflunomide<sup>36</sup>

Clinical diagnosis

**Approval Duration and Quantity Restrictions:**

**Approval:**

Initial Approval: Cytokine release syndrome = 1 month; members at high risk of developing ICANS = 3 months; immune checkpoint inhibitor-related toxicity = 6 months; all others = 12 months

Renewal Approval: Cytokine release syndrome = 1 month; members at high risk of developing ICANS = 3 months; immune checkpoint inhibitor-related toxicity = 6 months all others = 12 months

**Quantity Level Limit:**

- Kineret (anakinra) 100 mg/0.67 mL single-use prefilled syringe:



AETNA BETTER HEALTH®  
Coverage Policy/Guideline

Name: Kineret Page: 8 of 10

Effective Date: 2/2/2026 Last Review Date: 12/2025

Applies to:	<input type="checkbox"/> Illinois	<input type="checkbox"/> Florida	<input type="checkbox"/> Florida Kids
	<input checked="" type="checkbox"/> New Jersey	<input type="checkbox"/> Maryland	<input type="checkbox"/> Michigan
	<input type="checkbox"/> Pennsylvania Kids	<input type="checkbox"/> Virginia	<input type="checkbox"/> Kentucky PRMD

- 30 syringes per 30 days
- Exception limit: 360 syringes per 30 days

**References:**

1. Kineret [package insert]. Stockholm, Sweden: Swedish Orphan Biovitrum AB (publ); September 2024.
2. Micromedex® [electronic version]. Merative, Ann Arbor, Michigan, USA. Available at: <https://www.micromedexsolutions.com/>. Accessed April 18, 2025.
3. Ringold S, Weiss PF, Beukelman T, et al. 2013 Update of the 2011 American College of Rheumatology Recommendations for the Treatment of Juvenile Idiopathic Arthritis: Recommendations for the Medical Therapy of Children With Systemic Juvenile Idiopathic Arthritis and Tuberculosis Screening Among Children Receiving Biologic Medications. *Arthritis & Rheumatism*. 2013;65:2499-2512.
4. Laskari K, Tzioufas AG, Moutsopoulos HM. Efficacy and long-term follow-up of IL-1R inhibitor anakinra in adults with Still's disease: a case-series study. *Arthritis Res Ther*. 2011;13(3):R91.
5. Lequerre T, Quartier P, Rosellini D, et al. Interleukin-1 receptor antagonist (anakinra) treatment in patients with systemic-onset juvenile idiopathic arthritis or adult onset Still disease: preliminary experience in France. *Ann Rheum Dis*. 2008;67:302-308.
6. The NCCN Drugs & Biologics Compendium™. National Comprehensive Cancer Network, Inc. <http://www.nccn.org>. Accessed April 15, 2025.
7. Smolen JS, Landewé RBM, Bergstra SA, et al. EULAR recommendations for the management of rheumatoid arthritis with synthetic and biological disease-modifying antirheumatic drugs: 2022 update. *Ann Rheum Dis*. 2023;82:3-18.
8. Singh JA, Furst DE, Bharat A, et al. 2012 Update of the 2008 American College of Rheumatology recommendations for the use of disease-modifying antirheumatic drugs and biologic agents in the treatment of rheumatoid arthritis. *Arthritis Care Res*. 2012;64(5):625-639.
9. Saag KG, Teng GG, Patkar NM, et al. American College of Rheumatology 2008 recommendations for the use of nonbiologic and biologic disease-modifying antirheumatic drugs in rheumatoid arthritis. *Arthritis Rheum*. 2008;59(6):762-784.
10. Aletaha D, Neogi T, Silman AJ, et al. 2010 Rheumatoid Arthritis Classification Criteria. An American College of Rheumatology/European League Against Rheumatism Collaborative Initiative. *Arthritis Rheum*. 2010;62:2569-2581.
11. Anderson J, Caplan L, Yazdany J, et al. Rheumatoid Arthritis Disease Activity Measures: American College of Rheumatology Recommendations for Use in Clinical Practice. *Arthritis Rheum*. 2010;64:640-647.
12. Beukelman T, Patkar NM, Saag KG, et al. 2011 American College of Rheumatology recommendations for the treatment of juvenile idiopathic arthritis: initiation and safety monitoring



AETNA BETTER HEALTH®  
Coverage Policy/Guideline

Name: Kineret Page: 9 of 10

Effective Date: 2/2/2026 Last Review Date: 12/2025

Applies to:	<input type="checkbox"/> Illinois	<input type="checkbox"/> Florida	<input type="checkbox"/> Florida Kids
	<input checked="" type="checkbox"/> New Jersey	<input type="checkbox"/> Maryland	<input type="checkbox"/> Michigan
	<input type="checkbox"/> Pennsylvania Kids	<input type="checkbox"/> Virginia	<input type="checkbox"/> Kentucky PRMD

of therapeutic agents for the treatment of arthritis and systemic features. Arthritis Care Res. 2011;63(4):465-482.

13. Quartier P, Allantaz F, Cimaz R, et al. A multicentre, randomized, double-blind, placebo-controlled trial with the interleukin-1 receptor antagonist anakinra in patients with systemic-onset juvenile idiopathic arthritis (ANAJIS trial). *Ann Rheum Dis.* 2011;70:747-754.
14. Efthimiou P, Paik PK, Bielory L. Diagnosis and Management of Adult-Onset Still's Disease. *Ann Rheum Dis.* 2006;65(5):564-72. Epub 2005 Oct 11.
15. National Organization for Rare Disorders. Adult-Onset Still's Disease. URL: <https://rarediseases.org/rare-diseases/adult-onset-stills-disease>. Accessed November 18, 2024.
16. Adler Y, Charron P, Imazio M, et al. 2015 ESC Guidelines for the diagnosis and management of pericardial diseases: The Task Force for the Diagnosis and Management of Pericardial Diseases of the European Society of Cardiology (ESC) Endorsed by: The European Association for Cardio-Thoracic Surgery (EACTS). *Eur Heart J.* 2015;36(42):2921-64.
17. Kostjukovits S, Kalliokoski L, Antila K, et al. Treatment of hyperimmunoglobulinemia D Syndrome with biologics in children: review of the literature and Finnish experience. *Eur J Pediatr.* 2015;174(6):707-14.
18. National Organization for Rare Disorders. Mevalonate Kinase Deficiency. <http://rarediseases.org/rare-diseases/hyper-igd-syndrome>. Accessed November 18, 2024.
19. American College of Rheumatology. Hyperimmunoglobulin D Syndrome (HIDS)/Mevalonate Kinase Deficiency (MKD) (Juvenile). <https://rheumatology.org/patients/hyperimmunoglobulin-d-syndrome-hids-mevalonate-kinase-deficiency-mkd-juvenile>. Accessed November 18, 2024.
20. Simon A, Asli B, Braun-Falco M, et al. Schnitzler's syndrome: diagnosis, treatment, and follow-up. *Allergy.* 2013;68:562-568.
21. Lipsker D. The Schnitzler syndrome. *Orphanet J Rare Dis.* 2010;5:38.
22. Testing for TB Infection. Centers for Disease Control and Prevention. Retrieved on November 14, 2024 from: <https://www.cdc.gov/tb/testing/index.html>.
23. Lyseng-Williamson KA. Anakinra in Still's disease: A profile of its use. *Drugs Ther Perspect.* 2018;34(12):543-553.
24. Yoo DH. Biologics for the treatment of adult-onset still's disease. *Expert Opin Biol Ther.* 2019;19(11):1173-1190.
25. Richette P, Doherty M, Pascual E, et al. 2016 updated EULAR evidence-based recommendations for the management of gout. *Ann Rheum Dis.* 2017;76:29-42.
26. Zhang W, Doherty M, Pascual E, et al. EULAR recommendations for calcium pyrophosphate deposition. Part II: Management. *Ann Rheum Dis.* 2011;70:571-575.
27. Ottaviani S., Brunier L, Sibilía J, et al. Efficacy of anakinra in calcium pyrophosphate crystal-induced arthritis: A report of 16 cases and review of the literature. *Joint Bone Spine.* 2013;80:178-182.



AETNA BETTER HEALTH®  
Coverage Policy/Guideline

Name: Kineret Page: 10 of 10

Effective Date: 2/2/2026 Last Review Date: 12/2025

Applies to:	<input type="checkbox"/> Illinois	<input type="checkbox"/> Florida	<input type="checkbox"/> Florida Kids
	<input checked="" type="checkbox"/> New Jersey	<input type="checkbox"/> Maryland	<input type="checkbox"/> Michigan
	<input type="checkbox"/> Pennsylvania Kids	<input type="checkbox"/> Virginia	<input type="checkbox"/> Kentucky PRMD

28. FitzGerald JD, Dalbeth N, Mikuls T, et al. 2020 American College of Rheumatology Guideline for the Management of Gout [published correction appears in Arthritis Care Res (Hoboken). 2020;72(8):1187]. Arthritis Care Res (Hoboken). 2020;72(6):744-760.
29. Aksentijevich I, Masters SL, Ferguson PJ, et al. An autoinflammatory disease with deficiency of the interleukin-1-receptor antagonist. N Engl J Med. 2009;360(23):2426-37.
30. Bonilla FA, Khan DA, Ballas ZK, et al. Practice parameter for the diagnosis and management of primary immunodeficiency. J Allergy Clin Immunol. 2015;136(5):1186-205.
31. Smolen JS, Aletaha D. Assessment of rheumatoid arthritis disease activity and physical function. In: UpToDate, Post TW (Ed), UpToDate, Waltham, MA. Available with subscription. URL: www.uptodate.com. Accessed November 18, 2024.
32. Fraenkel L, Bathon JM, England BR, et al. 2021 American College of Rheumatology guideline for the treatment of rheumatoid arthritis. Arthritis Care Res. 2021;73(7):924-939.
33. Chiabrando JG, Bonaventura A, Vecchié A, et al. Management of acute and recurrent pericarditis: JACC State-of-the-art review. J Am Coll Cardiol. 2020;75(1):76-92.
34. Efthimiou P, Kontzias A, Hur P, et al. Adult-onset Still's disease in focus: Clinical manifestations, diagnosis, treatment, and unmet needs in the era of targeted therapies. Semin Arthritis Rheum. 2021;51(4):858-874.
35. Onel KB, Horton DB, Lovell DJ, et al. 2021 American College of Rheumatology guideline for the treatment of juvenile idiopathic arthritis: therapeutic approaches for oligoarthritis, temporomandibular joint arthritis, and systemic juvenile idiopathic arthritis. Arthritis Rheumatol. 2022;74(4):553-569.
36. Menter A, Gelfand JM, Connor C, et al. Joint American Academy of Dermatology-National Psoriasis Foundation guidelines of care for the management of psoriasis with systemic nonbiologic therapies. J Am Acad Dermatol. 2020;82(6):1445-1486.
37. Kumar S, Furqan M, Kafil T, et al. (2023). The paradigm shift in the management of recurrent pericarditis. American College of Cardiology. <https://www.acc.org/Latest-in-Cardiology/Articles/2022/12/19/14/52/The-Paradigm-Shift-in-the-Management-of-Recurrent-Pericarditis>.