



AETNA BETTER HEALTH®
Coverage Policy/Guideline

Name: Modafinil

Page: 1 of 4

Effective Date: 5/28/2025

Last Review Date: 5/2025

Applies Illinois

Florida

Michigan

to: New Jersey

Maryland

Florida Kids

Intent:

The intent of this policy/guideline is to provide information to the prescribing practitioner outlining the coverage criteria for modafinil under the patient's prescription drug benefit.

Description:

Modafinil) is indicated to improve wakefulness in adult patients with excessive sleepiness associated with narcolepsy, obstructive sleep apnea, or shift work disorder.

Limitations of Use

In obstructive sleep apnea (OSA), modafinil is indicated to treat excessive sleepiness and not as treatment for the underlying obstruction. If continuous positive airway pressure (CPAP) is the treatment of choice for a patient, a maximal effort to treat with CPAP for an adequate period of time should be made prior to initiating and during treatment with Provigil (modafinil) for excessive sleepiness.

Compensial Uses/Limited Treatment Option

Fatigue related to multiple sclerosis^{8,9}

Idiopathic hypersomnia⁶

Applicable Drug List:

Modafinil

Policy/Guideline:

Coverage Criteria

Idiopathic Hypersomnia

Authorization may be granted when the requested drug is being prescribed for idiopathic hypersomnia when ALL of the following criteria are met:

- The requested drug is being prescribed by, or in consultation with, a sleep specialist.
- The patient has experienced the presence of daytime lapses into sleep or daily irrepressible periods of need to sleep for at least 3 months.
- Insufficient sleep syndrome has been ruled out such as by lack of improvement of sleepiness after an adequate trial of increased nocturnal time in bed, preferably confirmed by at least a week of sleep log with wrist actigraphy.
- A multiple sleep latency test (MSLT) documented fewer than two sleep onset rapid eye movement periods (SOREMPs) or no SOREMPs if the rapid eye movement (REM) latency on the preceding polysomnogram was less than or equal to 15 minutes.



AETNA BETTER HEALTH®
Coverage Policy/Guideline

Name: Modafinil

Page: 2 of 4

Effective Date: 5/28/2025

Last Review Date: 5/2025

Applies Illinois

Florida

Michigan

to: New Jersey

Maryland

Florida Kids

- Sleep lab evaluation showed at least ONE of the following: mean sleep latency on MLST of less than or equal to 8 minutes, total 24-hour sleep time of greater than or equal to 660 minutes on 24-hour polysomnographic monitoring after correcting any chronic sleep deprivation or by wrist actigraphy in association with a sleep log and averaged over at least 7 days of unrestricted sleep.
- The patient does NOT have cataplexy.
- Hypersomnolence or MSLT results are not better explained by ANY of the following: another sleep disorder, other medical or psychiatric disorder, use of drugs or medications.

Multiple Sclerosis-Related Fatigue

Authorization may be granted when the requested drug is being prescribed for multiple sclerosis-related fatigue.

Narcolepsy

Authorization may be granted for a diagnosis of excessive sleepiness associated with narcolepsy when ALL of the following criteria are met:

- The requested drug is being prescribed by, or in consultation with, a sleep specialist.
- The diagnosis is confirmed by sleep study.

Obstructive Sleep Apnea (OSA)

Authorization may be granted for a diagnosis of excessive sleepiness associated with obstructive sleep apnea (OSA) when ALL of the following criteria are met:

- The requested drug is being prescribed by, or in consultation with, a sleep specialist.
- The diagnosis has been confirmed by polysomnography or home sleep apnea test (HSAT) with a technically adequate device.
- The patient has been receiving treatment for the underlying airway obstruction (continuous positive airway pressure [CPAP] or bilevel positive airway pressure [BIPAP]) for at least one month.
- The patient will continue to use CPAP or BIPAP after the requested drug is started.

Shift Work Disorder (SWD)

Authorization may be granted for a diagnosis of excessive sleepiness associated with shift work disorder (SWD) when ALL of the following criteria are met:

- The requested drug is being prescribed by, or in consultation with, a sleep specialist.



AETNA BETTER HEALTH®
Coverage Policy/Guideline

Name: Modafinil

Page: 3 of 4

Effective Date: 5/28/2025

Last Review Date: 5/2025

Applies Illinois

Florida

Michigan

to: New Jersey

Maryland

Florida Kids

- A sleep log and actigraphy monitoring have been completed for at least 14 days AND show a disrupted sleep and wake pattern.
- Symptoms have been present for 3 or more months.

Continuation of Therapy

Idiopathic Hypersomnia

Authorization may be granted when the requested drug is being prescribed for idiopathic hypersomnia when the following criteria is met:

- The patient has achieved or maintained a positive response to treatment from baseline.

Multiple Sclerosis-Related Fatigue

Authorization may be granted when the requested drug is being prescribed for multiple sclerosis-related fatigue when the following criteria is met:

- The patient has achieved or maintained a positive response to treatment from baseline.

Narcolepsy

Authorization may be granted for a diagnosis of excessive sleepiness associated with narcolepsy when the following criteria is met:

- The patient has achieved or maintained a positive response to treatment from baseline.

Obstructive Sleep Apnea (OSA)

Authorization may be granted for a diagnosis of excessive sleepiness associated with obstructive sleep apnea (OSA) when ALL of the following criteria are met:

- The patient has achieved or maintained a positive response to treatment from baseline.
- The patient is compliant with using continuous positive airway pressure (CPAP) or bilevel positive airway pressure (BIPAP).

Shift Work Disorder (SWD)

Authorization may be granted for a diagnosis of excessive sleepiness associated with shift work disorder (SWD) when ALL of the following criteria are met:



AETNA BETTER HEALTH®
Coverage Policy/Guideline

Name: Modafinil Page: 4 of 4

Effective Date: 5/28/2025 Last Review Date: 5/2025

Applies to: Illinois Florida Michigan
 New Jersey Maryland Florida Kids

- The patient has achieved or maintained a positive response to treatment from baseline.
- The patient is still a shift-worker.

Approval Duration and Quantity Restrictions:

Approval: 12 months

Quantity Level Limit: 60 tablets per 30 days

References:

1. Provigil [package insert]. Parsippany, NJ: Teva Pharmaceuticals; December 2022.
2. Lexicomp Online, AHFS DI (Adult and Pediatric) Online. Waltham, MA: UpToDate, Inc.; 2024. <https://online.lexi.com>. Accessed November 19, 2024.
3. Micromedex® (electronic version). Merative, Ann Arbor, Michigan, USA. Available at: <https://www.micromedexsolutions.com/> (cited: 11/19/2024).
4. Kapur VK, Auckley DH, Chowdhuri S, et al. Clinical Practice Guideline for Diagnostic Testing for Adult Obstructive Sleep Apnea: An American Academy of Sleep Medicine Clinical Practice Guideline. *J Clin Sleep Med*. 2017;13(3):479-504.
5. Epstein LJ, Kristo D, Strollo PJ, et al. Clinical Guidelines for the Evaluation, Management and Long-term Care of Obstructive Sleep Apnea in Adults. *J Clin Sleep Med*. 2009;5(3):263-276.
6. American Academy of Sleep Medicine. International Classification of Sleep Disorders, Third Edition, Text Revision. American Academy of Sleep Medicine, 2023.
7. Sateia MJ. International Classification of Sleep Disorders- Third Edition: Highlights and Modifications. *CHEST*. 2014;146(5):1387-1394.
8. Maski K, Trotti LM, Kotagal S, et al. Treatment of central disorders of hypersomnolence: an American Academy of Sleep Medicine clinical practice guideline. *J Clin Sleep Med*. 2021;17(9):1881-1893.
9. Maski K, Trotti LM, Kotagal S, et al. Treatment of central disorders of hypersomnolence: an American Academy of Sleep Medicine systematic review, meta-analysis, and GRADE assessment. *J Clin Sleep Med*. 2021;17(9):1895-1945.
10. National Multiple Sclerosis Society. Fatigue. Available at: <https://www.nationalmssociety.org/Symptoms-Diagnosis/MS-Symptoms/Fatigue>. Accessed November 19, 2024.
11. Shangyan H, Kuiqing L, Yumin X, et al. Meta-analysis of the efficiency of modafinil versus placebo in the treatment of multiple sclerosis fatigue. *Mult Scler Relat Disord*. 2018;19:85-89.
12. Trotti LM, Becker LA, Friederich Murray C, et al. Medications for daytime sleepiness in individuals with idiopathic hypersomnia. *Cochrane Database Syst Rev*. 2021;5(5):CD012714.