



AETNA BETTER HEALTH®
Coverage Policy/Guideline

Name: Neulasta and pegfilgrastim biosimilars Page: 1 of 9

Effective Date: 1/29/2024 Last Review Date: 12/2023

| | | | |
|-------------|----------------------------------------------|-----------------------------------|---------------------------------------|
| Applies to: | <input checked="" type="checkbox"/> Illinois | <input type="checkbox"/> Florida | <input type="checkbox"/> Florida Kids |
| | <input type="checkbox"/> New Jersey | <input type="checkbox"/> Maryland | <input type="checkbox"/> Michigan |
| | <input type="checkbox"/> Pennsylvania Kids | <input type="checkbox"/> Virginia | <input type="checkbox"/> Arizona |

Intent:

The intent of this policy/guideline is to provide information to the prescribing practitioner outlining the coverage criteria for Neulasta and pegfilgrastim biosimilars under the patient's prescription drug benefit.

Description:

The indications below including FDA-approved indications and compendial uses are considered a covered benefit provided that all the approval criteria are met and the member has no exclusions to the prescribed therapy.

A. FDA-Approved Indication

Neulasta

1. Patients with Cancer Receiving Myelosuppressive Chemotherapy
Neulasta is indicated to decrease the incidence of infection, as manifested by febrile neutropenia, in patients with non-myeloid malignancies receiving myelosuppressive anti-cancer drugs associated with a clinically significant incidence of febrile neutropenia.
2. Hematopoietic Subsyndrome of Acute Radiation Syndrome
Neulasta is indicated to increase survival in patients acutely exposed to myelosuppressive doses of radiation (Hematopoietic Subsyndrome of Acute Radiation Syndrome).

Fulphila²

Patients with Cancer Receiving Myelosuppressive Chemotherapy
Fulphila is indicated to decrease the incidence of infection, as manifested by febrile neutropenia, in patients with non-myeloid malignancies receiving myelosuppressive anti-cancer drugs associated with a clinically significant incidence of febrile neutropenia

Udenyca

Patients with Cancer Receiving Myelosuppressive Chemotherapy
Udenyca is indicated to decrease the incidence of infection, as manifested by febrile neutropenia, in patients with non-myeloid malignancies receiving myelosuppressive anti-cancer drugs associated with a clinically significant incidence of febrile neutropenia.

Ziextenzo

Patients with Cancer Receiving Myelosuppressive Chemotherapy



AETNA BETTER HEALTH®
Coverage Policy/Guideline

Name: Neulasta and pegfilgrastim biosimilars Page: 2 of 9

Effective Date: 1/29/2024 Last Review Date: 12/2023

| | | | |
|-------------|----------------------------------------------|-----------------------------------|---------------------------------------|
| Applies to: | <input checked="" type="checkbox"/> Illinois | <input type="checkbox"/> Florida | <input type="checkbox"/> Florida Kids |
| | <input type="checkbox"/> New Jersey | <input type="checkbox"/> Maryland | <input type="checkbox"/> Michigan |
| | <input type="checkbox"/> Pennsylvania Kids | <input type="checkbox"/> Virginia | <input type="checkbox"/> Arizona |

Ziextenzo is indicated to decrease the incidence of infection, as manifested by febrile neutropenia, in patients with non-myeloid malignancies receiving myelosuppressive anti-cancer drugs associated with a clinically significant incidence of febrile neutropenia.

Nyvepria

Patients with Cancer Receiving Myelosuppressive Chemotherapy

Nyvepria is indicated to decrease the incidence of infection, as manifested by febrile neutropenia, in patients with non-myeloid malignancies receiving myelosuppressive anti-cancer drugs associated with a clinically significant incidence of febrile neutropenia.

Fylnetra

Patients with Cancer Receiving Myelosuppressive Chemotherapy

Fylnetra is indicated to decrease the incidence of infection, as manifested by febrile neutropenia, in patients with non-myeloid malignancies receiving myelosuppressive anti-cancer drugs associated with a clinically significant incidence of febrile neutropenia.

Stimufend

Patients with Cancer Receiving Myelosuppressive Chemotherapy

Stimufend is indicated to decrease the incidence of infection, as manifested by febrile neutropenia, in patients with non-myeloid malignancies receiving myelosuppressive anti-cancer drugs associated with a clinically significant incidence of febrile neutropenia.

B. Compendial Use

1. Stem cell transplantation-related indications
2. Prophylaxis for chemotherapy-induced febrile neutropenia in patients with solid tumors
3. Hematopoietic Subsyndrome of Acute Radiation Syndrome
4. Hairy cell leukemia, neutropenic fever

All other indications are considered experimental/investigational and not medically necessary.



AETNA BETTER HEALTH®
Coverage Policy/Guideline

Name: Neulasta and pegfilgrastim biosimilars Page: 3 of 9

Effective Date: 1/29/2024 Last Review Date: 12/2023

| | | | |
|-------------|----------------------------------------------|-----------------------------------|---------------------------------------|
| Applies to: | <input checked="" type="checkbox"/> Illinois | <input type="checkbox"/> Florida | <input type="checkbox"/> Florida Kids |
| | <input type="checkbox"/> New Jersey | <input type="checkbox"/> Maryland | <input type="checkbox"/> Michigan |
| | <input type="checkbox"/> Pennsylvania Kids | <input type="checkbox"/> Virginia | <input type="checkbox"/> Arizona |

Applicable Drug List:

Neulasta
Fulphila
Fylnetra
Nyvepria
Stimufend
Udenyca
Ziextenzo

Policy/Guideline:

Documentation:

Primary Prophylaxis of Febrile Neutropenia

- A. Documentation must be provided of the member's diagnosis and chemotherapeutic regimen.
- B. If chemotherapeutic regimen has an intermediate risk of febrile neutropenia (10-19% [See Appendix B]), documentation must be provided outlining the patient's risk factors that confirm the member is at high risk for febrile neutropenia.

Criteria for Initial Approval:

A. Prevention of neutropenia in cancer patients receiving myelosuppressive chemotherapy

Authorization of 6 months may be granted for prevention of febrile neutropenia when all of the following criteria are met (1, 2, 3, and 4):

1. The requested medication will not be used in combination with other colony stimulating factors within any chemotherapy cycle.
2. The member will not receive chemotherapy at the same time as they receive radiation therapy.
3. The requested medication will not be administered with weekly chemotherapy regimens.
4. One of the following criteria is met (i or ii):
 - i. The requested medication will be used for primary prophylaxis in members with a solid tumor or non-myeloid malignancies who have received, are currently receiving, or will be receiving any of the following:
 - a. Myelosuppressive anti-cancer therapy that is expected to result in 20% or higher incidence of febrile neutropenia (FN) (See Appendix A).
 - b. Myelosuppressive anti-cancer therapy that is expected to result in 10 – 19% risk of FN (See Appendix B) and who are considered to be at high risk of FN because of bone marrow compromise, co-morbidities, or other patient specific risk factors (See Appendix C).



AETNA BETTER HEALTH®
Coverage Policy/Guideline

Name: Neulasta and pegfilgrastim biosimilars Page: 4 of 9

Effective Date: 1/29/2024 Last Review Date: 12/2023

| | | | |
|-------------|----------------------------------------------|-----------------------------------|---------------------------------------|
| Applies to: | <input checked="" type="checkbox"/> Illinois | <input type="checkbox"/> Florida | <input type="checkbox"/> Florida Kids |
| | <input type="checkbox"/> New Jersey | <input type="checkbox"/> Maryland | <input type="checkbox"/> Michigan |
| | <input type="checkbox"/> Pennsylvania Kids | <input type="checkbox"/> Virginia | <input type="checkbox"/> Arizona |

- c. Myelosuppressive anti-cancer therapy that is expected to result in less than 10% risk of FN and who have at least 2 patient-related risk factors (See *Appendix C*).
- ii. The requested medication will be used for secondary prophylaxis in members with solid tumors or non-myeloid malignancies who experienced a febrile neutropenic complication or a dose-limiting neutropenic event (a nadir or day of treatment count impacting the planned dose of chemotherapy) from a prior cycle of similar chemotherapy, with the same dose and scheduled planned for the current cycle (for which primary prophylaxis was not received).

B. Other indications

Authorization of 6 months may be granted for members with any of the following indications:

1. Stem cell transplantation-related indications
2. Hematopoietic Subsyndrome of Acute Radiation Syndrome
Treatment for radiation-induced myelosuppression following a radiological/nuclear incident
3. Hairy cell leukemia
Members with hairy cell leukemia with neutropenic fever following chemotherapy

Continuation of Therapy:

All members (including new members) requesting authorization for continuation of therapy must meet all initial authorization criteria.

Appendix:

A. APPENDIX A: Selected Chemotherapy Regimens with an Incidence of Febrile Neutropenia of 20% or Higher^{*†}

1. Acute Lymphoblastic Leukemia:
Select ALL regimens as directed by treatment protocol (see NCCN guidelines ALL)
2. Bladder Cancer:
 - i. Dose dense MVAC (methotrexate, vinblastine, doxorubicin, cisplatin)
 - ii. CBDCa/Pac (carboplatin, paclitaxel)
3. Bone Cancer
 - i. VAI (vincristine, doxorubicin or dactinomycin, ifosfamide)
 - ii. VDC-IE (vincristine, doxorubicin or dactinomycin, and cyclophosphamide alternating with ifosfamide and etoposide)
 - iii. Cisplatin/doxorubicin
 - iv. VDC (cyclophosphamide, vincristine, doxorubicin or dactinomycin)



AETNA BETTER HEALTH®
Coverage Policy/Guideline

Name: Neulasta and pegfilgrastim biosimilars Page: 5 of 9

Effective Date: 1/29/2024 Last Review Date: 12/2023

| | | | |
|-------------|----------------------------------------------|-----------------------------------|---------------------------------------|
| Applies to: | <input checked="" type="checkbox"/> Illinois | <input type="checkbox"/> Florida | <input type="checkbox"/> Florida Kids |
| | <input type="checkbox"/> New Jersey | <input type="checkbox"/> Maryland | <input type="checkbox"/> Michigan |
| | <input type="checkbox"/> Pennsylvania Kids | <input type="checkbox"/> Virginia | <input type="checkbox"/> Arizona |

- v. VIDE (vincristine, ifosfamide, doxorubicin or dactinomycin, etoposide)
- 4. Breast Cancer:
 - i. Docetaxel + trastuzumab
 - ii. Dose-dense AC (doxorubicin, cyclophosphamide) + paclitaxel (or dose dense paclitaxel)
 - iii. TAC (docetaxel, doxorubicin, cyclophosphamide)
 - iv. AT (doxorubicin, docetaxel)
 - v. Doc (docetaxel)
 - vi. TC (docetaxel, cyclophosphamide)
 - vii. TCH (docetaxel, carboplatin, trastuzumab)
- 5. Colorectal Cancer:
FOLFOXIRI (fluorouracil, leucovorin, oxaliplatin, irinotecan)
- 6. Esophageal and Gastric Cancers:
Docetaxel/cisplatin/fluorouracil
- 7. Head and Neck Squamous Cell Carcinoma
TPF (docetaxel, cisplatin, 5-fluorouracil)
- 8. Hodgkin Lymphoma:
 - i. Brentuximab vedotin + AVD (doxorubicin, vinblastine, dacarbazine)
 - ii. Escalated BEACOPP (bleomycin, etoposide, doxorubicin, cyclophosphamide, vincristine, procarbazine, prednisone)
- 9. Kidney Cancer:
Doxorubicin/gemcitabine
- 10. Non-Hodgkin's Lymphoma:
 - i. CHP (cyclophosphamide, doxorubicin, prednisone) + brentuximab vedotin
 - ii. Dose-adjusted EPOCH (etoposide, prednisone, vincristine, cyclophosphamide, doxorubicin)
 - iii. ICE (ifosfamide, carboplatin, etoposide)
 - iv. Dose-dense CHOP-14 (cyclophosphamide, doxorubicin, vincristine, prednisone) ± rituximab
 - v. MINE (mesna, ifosfamide, mitoxantrone, etoposide)
 - vi. DHAP (dexamethasone, cisplatin, cytarabine)
 - vii. ESHAP (etoposide, methylprednisolone, cisplatin, cytarabine (Ara-C))
 - viii. HyperCVAD ± rituximab (cyclophosphamide, vincristine, doxorubicin, dexamethasone ± rituximab)
 - ix. VAPEC-B (vincristine, doxorubicin, prednisolone, etoposide, cyclophosphamide, bleomycin)
- 11. Melanoma:
Dacarbazine-based combination with IL-2, interferon alpha (dacarbazine, cisplatin, vinblastine, IL-2, interferon alfa)



AETNA BETTER HEALTH®
Coverage Policy/Guideline

Name: Neulasta and pegfilgrastim biosimilars Page: 6 of 9

Effective Date: 1/29/2024 Last Review Date: 12/2023

| | | | |
|-------------|----------------------------------------------|-----------------------------------|---------------------------------------|
| Applies to: | <input checked="" type="checkbox"/> Illinois | <input type="checkbox"/> Florida | <input type="checkbox"/> Florida Kids |
| | <input type="checkbox"/> New Jersey | <input type="checkbox"/> Maryland | <input type="checkbox"/> Michigan |
| | <input type="checkbox"/> Pennsylvania Kids | <input type="checkbox"/> Virginia | <input type="checkbox"/> Arizona |

12. Multiple Myeloma:
 - i. VTD-PACE
(dexamethasone/thalidomide/cisplatin/doxorubicin/cyclophosphamide/etoposide + bortezomib)
 - ii. DT-PACE
(dexamethasone/thalidomide/cisplatin/doxorubicin/cyclophosphamide/etoposide)
13. Ovarian Cancer:
 - i. Topotecan
 - ii. Docetaxel
14. Pancreatic Cancer:
FOLFIRINOX (fluorouracil, leucovorin, irinotecan, oxaliplatin)
15. Soft Tissue Sarcoma:
 - i. MAID (mesna, doxorubicin, ifosfamide, dacarbazine)
 - ii. Doxorubicin
 - iii. Ifosfamide/doxorubicin
16. Small Cell Lung Cancer:
 - i. Top (topotecan)
 - ii. CAV (cyclophosphamide, doxorubicin, vincristine)
17. Testicular Cancer:
 - i. Velp (vinblastine, ifosfamide, cisplatin)
 - ii. VIP (etoposide, ifosfamide, cisplatin)
 - iii. TIP (paclitaxel, ifosfamide, cisplatin)
18. Gestational Trophoblastic Neoplasia:
 - i. EMA/EP (etoposide, methotrexate, dactinomycin/etoposide, cisplatin)
 - ii. EP/EMA (etoposide, cisplatin/etoposide, methotrexate, dactinomycin)
 - iii. TP/TE (paclitaxel, cisplatin/paclitaxel, etoposide)
 - iv. BEP (bleomycin, etoposide, cisplatin)
 - v. VIP (etoposide, ifosfamide, cisplatin)
 - vi. ICE (ifosfamide, carboplatin, etoposide)
19. Wilms Tumor:
 - i. Regimen M (vincristine, dactinomycin, doxorubicin, cyclophosphamide, etoposide)
 - ii. Regimen I (vincristine, doxorubicin, cyclophosphamide, etoposide)

*Applies to chemotherapy regimens with or without monoclonal antibodies (e.g., trastuzumab, rituximab)

† This list is not comprehensive; there are other agents/regimens that have an intermediate/high risk for development of febrile neutropenia.



AETNA BETTER HEALTH®
Coverage Policy/Guideline

Name: Neulasta and pegfilgrastim biosimilars Page: 7 of 9

Effective Date: 1/29/2024 Last Review Date: 12/2023

| | | | |
|-------------|----------------------------------------------|-----------------------------------|---------------------------------------|
| Applies to: | <input checked="" type="checkbox"/> Illinois | <input type="checkbox"/> Florida | <input type="checkbox"/> Florida Kids |
| | <input type="checkbox"/> New Jersey | <input type="checkbox"/> Maryland | <input type="checkbox"/> Michigan |
| | <input type="checkbox"/> Pennsylvania Kids | <input type="checkbox"/> Virginia | <input type="checkbox"/> Arizona |

B. APPENDIX B: Selected Chemotherapy Regimens with an Incidence of Febrile Neutropenia of 10% to 19%*†

1. Occult Primary – Adenocarcinoma:
Gemcitabine/docetaxel
2. Breast Cancer:
 - i. Docetaxel
 - ii. CMF classic (cyclophosphamide, methotrexate, fluorouracil)
 - iii. CA (doxorubicin, cyclophosphamide) (60 mg/m²) (hospitalized)
 - iv. AC (doxorubicin, cyclophosphamide) + sequential docetaxel (taxane portion only)
 - v. AC + sequential docetaxel + trastuzumab
 - vi. A (doxorubicin) (75 mg/m²)
 - vii. AC (doxorubicin, cyclophosphamide)
 - viii. CapDoc (capecitabine, docetaxel)
 - ix. Paclitaxel every 21 days
3. Cervical Cancer:
 - i. Irinotecan
 - ii. Cisplatin/topotecan
 - iii. Paclitaxel/cisplatin
 - iv. Topotecan
4. Colorectal Cancer:
 - i. FL (fluorouracil, leucovorin)
 - ii. CPT-11 (irinotecan) (350 mg/m² q 3 wk)
 - iii. FOLFOX (fluorouracil, leucovorin, oxaliplatin)
5. Esophageal and Gastric Cancers:
 - i. Irinotecan/cisplatin
 - ii. Epirubicin/cisplatin/5-fluorouracil
 - iii. Epirubicin/cisplatin/capecitabine
6. Non-Hodgkin's Lymphomas:
 - i. EPOCH-IT chemotherapy
 - ii. GDP (gemcitabine, dexamethasone, cisplatin/carboplatin)
 - iii. GDP (gemcitabine, dexamethasone, cisplatin/carboplatin) + rituximab
 - iv. FMR (fludarabine, mitoxantrone, rituximab)
 - v. CHOP (cyclophosphamide, doxorubicin, vincristine, prednisone) including regimens with pegylated liposomal doxorubicin
 - vi. CHOP + rituximab (cyclophosphamide, doxorubicin, vincristine, prednisone, rituximab) including regimens with pegylated liposomal doxorubicin
 - vii. Bendamustine
7. Non-Small Cell Lung Cancer:
 - i. Cisplatin/paclitaxel



AETNA BETTER HEALTH®
Coverage Policy/Guideline

Name: Neulasta and pegfilgrastim biosimilars Page: 8 of 9

Effective Date: 1/29/2024 Last Review Date: 12/2023

| | | | |
|-------------|----------------------------------------------|-----------------------------------|---------------------------------------|
| Applies to: | <input checked="" type="checkbox"/> Illinois | <input type="checkbox"/> Florida | <input type="checkbox"/> Florida Kids |
| | <input type="checkbox"/> New Jersey | <input type="checkbox"/> Maryland | <input type="checkbox"/> Michigan |
| | <input type="checkbox"/> Pennsylvania Kids | <input type="checkbox"/> Virginia | <input type="checkbox"/> Arizona |

- ii. Cisplatin/vinorelbine
- iii. Cisplatin/docetaxel
- iv. Cisplatin/etoposide
- v. Carboplatin/paclitaxel
- vi. Docetaxel
- 8. Ovarian Cancer:
Carboplatin/docetaxel
- 9. Prostate Cancer:
Cabazitaxel
- 10. Small Cell Lung Cancer:
Etoposide/carboplatin
- 11. Testicular Cancer:
 - i. BEP (bleomycin, etoposide, cisplatin)
 - ii. Etoposide/cisplatin
- 12. Uterine Sarcoma:
Docetaxel

*Applies to chemotherapy regimens with or without monoclonal antibodies (e.g., trastuzumab, rituximab)

† This list is not comprehensive; there are other agents/regimens that have an intermediate/high risk for development of febrile neutropenia.

Approval Duration and Quantity Restrictions:

Approval: 6 months

Quantity Level Limit: Neulasta/Fulphila/Fylnetra/Nyvepria/Stimufend/ Udenyca/Ziextenzo (pegfilgrastim) injection 6 mg per 0.6 mL solution: 2 per 28 days

References:

1. Neulasta [package insert]. Thousand Oaks, CA: Amgen Inc.; February 2021.
2. Fulphila [package insert]. Morgantown, WV: Mylan Pharmaceuticals, Inc.; October 2021.
3. Udenyca [package insert]. Redwood City, CA: Coherus BioSciences, Inc.; March 2023.
4. Ziextenzo [package insert]. Princeton, NJ: Sandoz Inc.; March 2021.
5. Nyvepria [package insert]. Lake Forest, IL: Hospira, Inc.; March 2023.
6. Fylnetra [package insert]. Piscataway, NJ: Kashiv BioSciences, LLC; May 2022.
7. Stimufend [package insert]. Lake Zurich, IL: Fresenius Kabi USA, LLC; September 2022.
8. The NCCN Drugs & Biologics Compendium® © 2023 National Comprehensive Cancer Network, Inc. Available at: <https://www.nccn.org> Accessed June 20, 2023.
9. National Comprehensive Cancer Network. NCCN Clinical Practice Guidelines in Oncology: Hematopoietic Growth Factors. Version 2.2023.



AETNA BETTER HEALTH®
Coverage Policy/Guideline

Name: Neulasta and pegfilgrastim biosimilars Page: 9 of 9

Effective Date: 1/29/2024 Last Review Date: 12/2023

| | | | |
|-------------|----------------------------------------------|-----------------------------------|---------------------------------------|
| Applies to: | <input checked="" type="checkbox"/> Illinois | <input type="checkbox"/> Florida | <input type="checkbox"/> Florida Kids |
| | <input type="checkbox"/> New Jersey | <input type="checkbox"/> Maryland | <input type="checkbox"/> Michigan |
| | <input type="checkbox"/> Pennsylvania Kids | <input type="checkbox"/> Virginia | <input type="checkbox"/> Arizona |

https://www.nccn.org/professionals/physician_gls/pdf/growthfactors.pdf Accessed June 13, 2023.

10. IBM Micromedex® DRUGDEX® (electronic version). IBM Watson Health, Greenwood Village, Colorado, USA. Available at <https://www.micromedexsolutions.com> (Accessed: June 20, 2023).
11. Smith TJ, Bohlke K, Lyman GH, et al. Recommendations for the use of white blood cell growth factors: American Society of Clinical Oncology Clinical Practice Guideline Update. *J Clin Oncol*. 2015;33(28):3199-3212.
12. National Comprehensive Cancer Network. NCCN Clinical Practice Guidelines in Oncology: Hairy Cell Leukemia. Version 1.2023.
https://www.nccn.org/professionals/physician_gls/pdf/hairy_cell.pdf Accessed June 19, 2023.
13. Smith TJ, Khatcheressian J, Lyman GH, et al. 2006 update of recommendations for the use of white blood cell growth factors: an evidence-based clinical practice guideline. *J Clin Oncol*. 2006;24(19):3187-3205.
14. National Comprehensive Cancer Network. NCCN Clinical Practice Guidelines in Oncology: Gestational Trophoblastic Neoplasia. Version 1.2023.
https://www.nccn.org/professionals/physician_gls/pdf/gtn.pdf Accessed June 19, 2023.
15. National Comprehensive Cancer Network. NCCN Clinical Practice Guidelines in Oncology: Wilms Tumor (Nephroblastoma). Version 1.2023.
https://www.nccn.org/professionals/physician_gls/pdf/wilms_tumor.pdf Accessed June 19, 2023.