



AETNA BETTER HEALTH®
Coverage Policy/Guideline

Name: Simponi

Page: 1 of 10

Effective Date: 1/13/2026

Last Review Date: 12/2025

Applies to:	☒ Illinois	☐ Florida	☐ Florida Kids
	☐ New Jersey	☐ Maryland	☐ Michigan
	☐ Pennsylvania Kids	☐ Virginia	☐ Kentucky PRMD

Intent:

The intent of this policy/guideline is to provide information to the prescribing practitioner outlining the coverage criteria for Simponi under the patient's prescription drug benefit.

Description:

The indications below including FDA-approved indications and compendial uses are considered a covered benefit provided that all the approval criteria are met and the member has no exclusions to the prescribed therapy.

FDA-approved Indications¹

- Moderately to severely active rheumatoid arthritis (RA) in adults, in combination with methotrexate
- Active psoriatic arthritis (PsA) in adults, alone or in combination with methotrexate
- Active ankylosing spondylitis (AS) in adults
- Moderately to severely active ulcerative colitis (UC) in adults and pediatric patients weighing at least 15 kilograms (kg)

Compendial Uses

- Non-radiographic axial spondyloarthritis²
- Immune checkpoint inhibitor-related toxicities - inflammatory arthritis²¹

All other indications are considered experimental/investigational and not medically necessary.

Applicable Drug List:

Simponi

Policy/Guideline:

Documentation for all indications:

The patient is unable to take THREE preferred products, where indicated, for the given diagnosis due to a trial and inadequate treatment response or intolerance, or a contraindication. Documentation is required for approval.



AETNA BETTER HEALTH®
Coverage Policy/Guideline

Name: Simponi

Page: 2 of 10

Effective Date: 1/13/2026

Last Review Date: 12/2025

Applies to:	☒ Illinois	☐ Florida	☐ Florida Kids
	☐ New Jersey	☐ Maryland	☐ Michigan
	☐ Pennsylvania Kids	☐ Virginia	☐ Kentucky PRMD

Documentation

Submission of the following information is necessary to initiate the prior authorization review:

Rheumatoid Arthritis (RA)

Initial Requests

- Chart notes, medical record documentation, or claims history supporting previous medications tried (if applicable), including response to therapy. If therapy is not advisable, documentation of clinical reason to avoid therapy.
- Laboratory results, chart notes, or medical record documentation of biomarker testing (i.e., rheumatoid factor [RF], anti-cyclic citrullinated peptide [anti-CCP], and C-reactive protein [CRP] and/or erythrocyte sedimentation rate [ESR]) (if applicable).

Continuation Requests

Chart notes or medical record documentation supporting positive clinical response.

Ankylosing Spondylitis (AS), Non-Radiographic Axial Spondyloarthritis (nr-axSpA), Psoriatic Arthritis (PsA), and Immune Checkpoint Inhibitor-Related Toxicity

Initial Requests

Chart notes, medical record documentation, or claims history supporting previous medications tried (if applicable), including response to therapy. If therapy is not advisable, documentation of clinical reason to avoid therapy.

Continuation Requests

Chart notes or medical record documentation supporting positive clinical response.

Ulcerative Colitis (UC)

Continuation Requests

Chart notes or medical record documentation supporting positive clinical response to therapy or remission.

Prescriber Specialties

This medication must be prescribed by or in consultation with one of the following:

- Rheumatoid arthritis, ankylosing spondylitis, and non-radiographic axial spondyloarthritis: rheumatologist
- Psoriatic arthritis: rheumatologist or dermatologist
- Ulcerative colitis: gastroenterologist



AETNA BETTER HEALTH®
Coverage Policy/Guideline

Name: Simponi

Page: 3 of 10

Effective Date: 1/13/2026

Last Review Date: 12/2025

Applies to:	☒ Illinois	☐ Florida	☐ Florida Kids
	☐ New Jersey	☐ Maryland	☐ Michigan
	☐ Pennsylvania Kids	☐ Virginia	☐ Kentucky PRMD

- Immune checkpoint inhibitor-related toxicity: oncologist, hematologist, or rheumatologist

Coverage Criteria

Rheumatoid Arthritis (RA)^{1,3-5,16,18,19}

Authorization of 12 months may be granted for adult members who have previously received a biologic or targeted synthetic drug indicated for moderately to severely active rheumatoid arthritis within the past 120 days. The requested medication must be prescribed in combination with methotrexate or leflunomide unless the member has a clinical reason not to use methotrexate or leflunomide (see Appendix).

Authorization of 12 months may be granted for adult members for treatment of moderately to severely active RA when all of the following criteria are met:

- Member meets either of the following:
 - Member has been tested for either of the following biomarkers and the test was positive:
 - Rheumatoid factor (RF)
 - Anti-cyclic citrullinated peptide (anti-CCP)
 - Member has been tested for ALL of the following biomarkers:
 - RF
 - Anti-CCP
 - C-reactive protein (CRP) and/or erythrocyte sedimentation rate (ESR)
- Member meets ONE of the following:
 - Member has failed to achieve a low disease activity after a 3-month trial of methotrexate (MTX) monotherapy at a maximum titrated dose of at least 15 mg per week and meets any of the following conditions:
 - Member has had a documented inadequate response to MTX in combination with at least one other conventional synthetic drug (i.e., hydroxychloroquine and/or sulfasalazine) after a 3-month trial at a maximum tolerated dose(s).
 - Member has experienced a documented intolerable adverse event to hydroxychloroquine or sulfasalazine.
 - Member has a documented contraindication to hydroxychloroquine (see Appendix) and sulfasalazine (e.g., porphyria, intestinal or urinary obstruction).
 - Member has moderate to high disease activity.



AETNA BETTER HEALTH®
Coverage Policy/Guideline

Name: Simponi

Page: 4 of 10

Effective Date: 1/13/2026

Last Review Date: 12/2025

Applies to:	☒ Illinois	☐ Florida	☐ Florida Kids
	☐ New Jersey	☐ Maryland	☐ Michigan
	☐ Pennsylvania Kids	☐ Virginia	☐ Kentucky PRMD

- Member was unable to tolerate a 3-month trial of MTX monotherapy at a maximum titrated dose of at least 15 mg per week and meets any of the following conditions:
 - Member has had a documented inadequate response to MTX in combination with at least one other conventional synthetic drug (i.e., hydroxychloroquine and/or sulfasalazine) after a 3-month trial at a maximum tolerated dose(s).
 - Member has stopped taking MTX and has had a documented inadequate response to another conventional synthetic drug (i.e., leflunomide, hydroxychloroquine, and/or sulfasalazine) alone or in combination after a 3-month trial at a maximum tolerated dose(s).
 - Member has experienced a documented intolerable adverse event to leflunomide, hydroxychloroquine, or sulfasalazine.
 - Member has a documented contraindication to leflunomide, hydroxychloroquine (see Appendix), and sulfasalazine (e.g., porphyria, intestinal or urinary obstruction).
 - Member has moderate to high disease activity.
- Member has experienced a documented intolerable adverse event or has a documented contraindication to MTX (see Appendix), discontinues MTX, and meets any of the following conditions:
 - Member has had a documented inadequate response to another conventional synthetic drug (i.e., leflunomide, hydroxychloroquine, and/or sulfasalazine) alone or in combination after a 3-month trial at a maximum tolerated dose(s).
 - Member has experienced a documented intolerable adverse event to leflunomide, hydroxychloroquine, or sulfasalazine.
 - Member has a documented contraindication to leflunomide, hydroxychloroquine (see Appendix), and sulfasalazine (e.g., porphyria, intestinal or urinary obstruction).
 - Member has moderate to high disease activity.
- Member is prescribed the requested medication in combination with methotrexate or leflunomide or has a clinical reason not to use methotrexate or leflunomide (see Appendix).

Psoriatic Arthritis (PsA)^{1,6-9,14}

Authorization of 12 months may be granted for adult members who have previously received a biologic or targeted synthetic drug indicated for active psoriatic arthritis.

Authorization of 12 months may be granted for adult members for treatment of active psoriatic arthritis when either of the following criteria is met:

- Member has mild to moderate disease and meets one of the following criteria:



AETNA BETTER HEALTH®
Coverage Policy/Guideline

Name: Simponi

Page: 5 of 10

Effective Date: 1/13/2026

Last Review Date: 12/2025

Applies to:	☒ Illinois	☐ Florida	☐ Florida Kids
	☐ New Jersey	☐ Maryland	☐ Michigan
	☐ Pennsylvania Kids	☐ Virginia	☐ Kentucky PRMD

- Member has had an inadequate response to methotrexate, leflunomide, or another conventional synthetic drug (e.g., sulfasalazine) administered at an adequate dose and duration.
- Member has an intolerance or contraindication to methotrexate or leflunomide (see Appendix), or another conventional synthetic drug (e.g., sulfasalazine).
- Member has enthesitis or predominantly axial disease.
- Member has severe disease.

Ankylosing Spondylitis (AS) And Non-Radiographic Axial Spondyloarthritis (nr-axSpA)^{1,2,10,11}

Authorization of 12 months may be granted for adult members who have previously received a biologic or targeted synthetic drug indicated for active ankylosing spondylitis or active non-radiographic axial spondyloarthritis.

Authorization of 12 months may be granted for adult members for treatment of active ankylosing spondylitis or active non-radiographic axial spondyloarthritis when either of the following criteria is met:

- Member has experienced an inadequate response to at least two non-steroidal anti-inflammatory drugs (NSAIDs).
- Member has an intolerance or contraindication to two or more NSAIDs.

Ulcerative Colitis (UC)^{1,12,15,17}

Authorization of 12 months may be granted for treatment of moderately to severely active ulcerative colitis.

Immune Checkpoint Inhibitor-Related Toxicity²¹

Authorization of 12 months may be granted for treatment of immune checkpoint inhibitor-related toxicity when the member has moderate or severe immunotherapy-related inflammatory arthritis and meets either of the following:

- Member has had an inadequate response to corticosteroids or a conventional synthetic drug (e.g., methotrexate, sulfasalazine, leflunomide, hydroxychloroquine).
- Member has an intolerance or contraindication to corticosteroids and a conventional synthetic drug (e.g., methotrexate, sulfasalazine, leflunomide, hydroxychloroquine).



AETNA BETTER HEALTH®
Coverage Policy/Guideline

Name: Simponi

Page: 6 of 10

Effective Date: 1/13/2026

Last Review Date: 12/2025

Applies to:	☒ Illinois	☐ Florida	☐ Florida Kids
	☐ New Jersey	☐ Maryland	☐ Michigan
	☐ Pennsylvania Kids	☐ Virginia	☐ Kentucky PRMD

Continuation of Therapy

Rheumatoid Arthritis (RA)^{1,3-5,16,18,19}

Authorization of 12 months may be granted for all adult members (including new members) who are using the requested medication for moderately to severely active rheumatoid arthritis and who achieve or maintain a positive clinical response as evidenced by disease activity improvement of at least 20% from baseline in tender joint count, swollen joint count, pain, or disability.

Psoriatic Arthritis (PsA)^{1,6-9,14}

Authorization of 12 months may be granted for all adult members (including new members) who are using the requested medication for psoriatic arthritis and who achieve or maintain a positive clinical response as evidenced by low disease activity or improvement in signs and symptoms of the condition when there is improvement in any of the following from baseline:

- Number of swollen joints
- Number of tender joints
- Dactylitis
- Enthesitis
- Axial disease
- Skin and/or nail involvement
- Functional status
- C-reactive protein (CRP)

Ankylosing Spondylitis (AS) and Non-Radiographic Axial Spondyloarthritis (nr-axSpA)^{1,2,10,11}

Authorization of 12 months may be granted for all adult members (including new members) who are using the requested medication for ankylosing spondylitis or non-radiographic axial spondyloarthritis and who achieve or maintain a positive clinical response as evidenced by low disease activity or improvement in signs and symptoms of the condition when there is improvement in any of the following from baseline:

- Functional status
- Total spinal pain
- Inflammation (e.g., morning stiffness)
- Swollen joints
- Tender joints
- C-reactive protein (CRP)

Ulcerative Colitis (UC)^{1,12,15,17}

Authorization of 12 months may be granted for all members (including new members) who are using the requested medication for moderately to severely active ulcerative colitis and who achieve or maintain remission.



AETNA BETTER HEALTH®
Coverage Policy/Guideline

Name: Simponi

Page: 7 of 10

Effective Date: 1/13/2026

Last Review Date: 12/2025

Applies to:	☒ Illinois	☐ Florida	☐ Florida Kids
	☐ New Jersey	☐ Maryland	☐ Michigan
	☐ Pennsylvania Kids	☐ Virginia	☐ Kentucky PRMD

Authorization of 12 months may be granted for all members (including new members) who are using the requested medication for moderately to severely active ulcerative colitis and who achieve or maintain a positive clinical response as evidenced by low disease activity or improvement in signs and symptoms of the condition when there is improvement in any of the following from baseline:

- Stool frequency
- Rectal bleeding
- Urgency of defecation
- C-reactive protein (CRP)
- Fecal calprotectin (FC)
- Appearance of the mucosa on endoscopy, computed tomography enterography (CTE), magnetic resonance enterography (MRE), or intestinal ultrasound
- Improvement on a disease activity scoring tool (e.g., Ulcerative Colitis Endoscopic Index of Severity [UCEIS], Mayo score)

Immune Checkpoint Inhibitor-Related Toxicity

Authorization of 12 months may be granted for all members (including new members) who are using the requested medication for immunotherapy-related inflammatory arthritis and who achieve or maintain a positive clinical response with the requested medication as evidenced by low disease activity or improvement in signs and symptoms of the condition.

Other^{1,13}

For all indications: Member has had a documented negative tuberculosis (TB) test (which can include a tuberculosis skin test [TST] or an interferon-release assay [IGRA]) within 12 months of initiating therapy for persons who are naïve to biologic drugs or targeted synthetic drugs associated with an increased risk of TB.

If the screening testing for TB is positive, there must be further testing to confirm there is no active disease (e.g., chest x-ray). Do not administer the requested medication to members with active TB infection. If there is latent disease, TB treatment must be started before initiation of the requested medication.

For all indications: Member cannot use the requested medication concomitantly with any other biologic drug or targeted synthetic drug for the same indication.

Dosage and Administration

Approvals may be subject to dosing limits in accordance with FDA-approved labeling, accepted compendia, and/or evidence-based practice guidelines.



AETNA BETTER HEALTH®
Coverage Policy/Guideline

Name: Simponi

Page: 8 of 10

Effective Date: 1/13/2026

Last Review Date: 12/2025

Applies to:	☒ Illinois	☐ Florida	☐ Florida Kids
	☐ New Jersey	☐ Maryland	☐ Michigan
	☐ Pennsylvania Kids	☐ Virginia	☐ Kentucky PRMD

Appendix

Examples of Clinical Reasons to Avoid Pharmacologic Treatment with Methotrexate, Hydroxychloroquine, or Leflunomide²⁰

- Clinical diagnosis of alcohol use disorder, alcoholic liver disease, or other chronic liver disease
- Drug interaction
- Risk of treatment-related toxicity
- Pregnancy or currently planning pregnancy
- Breastfeeding
- Significant comorbidity prohibits use of systemic agents (e.g., liver or kidney disease, blood dyscrasias, uncontrolled hypertension)
- Hypersensitivity
- History of intolerance or adverse event

Approval Duration and Quantity Restrictions:

Approval: Initial and Renewal Approval: 12 months

Quantity Level Limit:

- Simponi (golimumab) 50 mg per 0.5 mL single-dose pre-filled syringe/autoinjector:
 - 1 syringe/ autoinjector per 28 days
- Simponi (golimumab) 100 mg per 1 mL single-dose pre-filled syringe/autoinjector:
 - 1 syringe/ autoinjector per 28 days
 - Exception limit: 3 syringes/ autoinjectors per 14 days

References:

1. Simponi [package insert]. Horsham, PA: Janssen Biotech, Inc.; October 2025.
2. van der Heijde D, Ramiro S, Landewe R, et al. 2016 Update of the international ASAS-EULAR management recommendations for axial spondyloarthritis. *Ann Rheum Dis*. 2017;76(6):978-991.
3. Smolen JS, Landewé RBM, Bergstra SA, et al. EULAR recommendations for the management of rheumatoid arthritis with synthetic and biological disease-modifying antirheumatic drugs: 2022 update. *Ann Rheum Dis*. 2023;82:3-18.
4. Singh JA, Saag KG, Bridges SL Jr, et al. 2015 American College of Rheumatology Guideline for the Treatment of Rheumatoid Arthritis. *Arthritis Rheumatol*. 2016;68(1):1-26.
5. Saag KG, Teng GG, Patkar NM, et al. American College of Rheumatology 2008 recommendations for the use of nonbiologic and biologic disease-modifying antirheumatic drugs in rheumatoid arthritis. *Arthritis Rheum*. 2008;59(6):762-784.
6. Menter A, Korman NJ, Elmets CA, et al. Guidelines of care for the management of psoriasis and psoriatic arthritis. Section 6: Guidelines of care for the treatment of psoriasis and psoriatic



AETNA BETTER HEALTH®
Coverage Policy/Guideline

Name: Simponi

Page: 9 of 10

Effective Date: 1/13/2026

Last Review Date: 12/2025

Applies to:	☒ Illinois	☐ Florida	☐ Florida Kids
	☐ New Jersey	☐ Maryland	☐ Michigan
	☐ Pennsylvania Kids	☐ Virginia	☐ Kentucky PRMD

arthritis: case-based presentations and evidence-based conclusions. J Am Acad Dermatol. 2011;65(1):137-174.

7. Gossec L, Baraliakos X, Kerschbaumer A, et al. European League Against Rheumatism (EULAR) recommendations for the management of psoriatic arthritis with pharmacological therapies; 2019 update. Ann Rheum Dis. 2020;79(6):700-712.
8. Gladman DD, Antoni C, Mease P, et al. Psoriatic arthritis: epidemiology, clinical features, course, and outcome. Ann Rheum Dis 2005;64(Suppl II):ii14-ii17.
9. Coates LC, Soriano ER, Corp N, et al. Group for Research and Assessment of Psoriasis and Psoriatic Arthritis (GRAPPA): updated treatment recommendations for psoriatic arthritis 2021. Nat Rev Rheumatol. 2022;18(8):465-479.
10. Braun J, van den Berg R, Baraliakos X, et al. 2010 update of the ASAS/EULAR recommendations for the management of ankylosing spondylitis. Ann Rheum Dis. 2011;70:896-904.
11. Ward MM, Deodhar A, Gensler LS, et al. 2019 Update of the American College of Rheumatology/Spondylitis Association of America/Spondyloarthritis Research and Treatment Network Recommendations for the Treatment of Ankylosing Spondylitis and Nonradiographic Axial Spondyloarthritis. Arthritis Rheumatol. 2019;71(10):1599-1613. doi:10.1002/art.41042.
12. Talley NJ, Abreu MT, Achkar J, et al. An evidence-based systematic review on medical therapies for inflammatory bowel disease. Am J Gastroenterol. 2011;106(Suppl 1):S2-S25.
13. Testing for TB Infection. Centers for Disease Control and Prevention. Retrieved on June 16, 2025 from: <https://www.cdc.gov/tb/testing/index.html>.
14. Singh JA, Guyatt G, Ogdie A, et al. 2018 American College of Rheumatology/National Psoriasis Foundation Guideline for the Treatment of Psoriatic Arthritis. Arthritis Rheumatol. 2019;71(1):5-32. doi:10.1002/art.40726.
15. Singh S, Loftus EV Jr, Limketkai BN, et al. AGA Living Clinical Practice Guideline on Pharmacological Management of Moderate-to-Severe Ulcerative Colitis. Gastroenterology. 2024;167(7):1307-1343.
16. Aletaha D, Neogi T, Silman AJ, et al. 2010 Rheumatoid arthritis classification criteria: an American College of Rheumatology/European League Against Rheumatism collaborative initiative. Arthritis Rheum. 2010;62(9):2569-81.
17. Rubin DT, Ananthakrishnan AN, Siegel CA, et al. ACG Clinical Guideline Update: Ulcerative Colitis in Adults. Am J Gastroenterol. 2025;120(6):1187-1224.
18. Smolen JS, Aletaha D. Assessment of rheumatoid arthritis activity in clinical trials and clinical practice. In: UpToDate, Post TW (Ed), UpToDate, Waltham, MA. Available with subscription. URL: www.uptodate.com. Accessed March 19, 2021.
19. Fraenkel L, Bathon JM, England BR, et al. 2021 American College of Rheumatology guideline for the treatment of rheumatoid arthritis. Arthritis Care Res. 2021;0:1-16.
20. Menter A, Gelfand JM, Connor C, et al. Joint AAD-NPF guidelines of care for the management of psoriasis with systemic nonbiologic therapies. J Am Acad Dermatol. 2020;82(6):1445-86.

AETNA BETTER HEALTH®
Coverage Policy/Guideline



Name: Simponi

Page: 10 of 10

Effective Date: 1/13/2026

Last Review Date: 12/2025

Applies
to:

☒ Illinois

☐ New Jersey

☐ Pennsylvania Kids

☐ Florida

☐ Maryland

☐ Virginia

☐ Florida Kids

☐ Michigan

☐ Kentucky PRMD

21. The NCCN Drugs & Biologics Compendium® © 2025 National Comprehensive Cancer Network, Inc. Available at: <http://www.nccn.org>. Accessed June 16, 2025.