



AETNA BETTER HEALTH®
Coverage Policy/Guideline

Name: Treprostini Page: 1 of 4

Effective Date: 10/15/2025 Last Review Date: 9/2025

| | | | |
|-------------|---|--|---|
| Applies to: | <input checked="" type="checkbox"/> Illinois | <input type="checkbox"/> Florida | <input checked="" type="checkbox"/> Florida Kids |
| | <input type="checkbox"/> New Jersey | <input checked="" type="checkbox"/> Maryland | <input type="checkbox"/> Michigan |
| | <input checked="" type="checkbox"/> Pennsylvania Kids | <input type="checkbox"/> Virginia | <input checked="" type="checkbox"/> Kentucky PRMD |

Intent:

The intent of this policy/guideline is to provide information to the prescribing practitioner outlining the coverage criteria for treprostini under the patient's prescription drug benefit.

Description:

Indications

The indications below including FDA-approved indications and compendial uses are considered a covered benefit provided that all the approval criteria are met and the member has no exclusions to the prescribed therapy.

FDA-approved Indications^{1,2}

- Treatment of pulmonary arterial hypertension (PAH; World Health Organization [WHO] Group 1) to diminish symptoms associated with exercise. Studies establishing effectiveness included patients with New York Heart Association (NYHA) Functional Class II-IV symptoms and etiologies of idiopathic or heritable PAH, PAH associated with congenital systemic-to-pulmonary shunts, or PAH associated with connective tissue diseases.
- In patients with PAH requiring transition from epoprostenol, to diminish the rate of clinical deterioration. Consider the risks and benefits of each drug prior to transition.

All other indications are considered experimental/investigational and not medically necessary.

Applicable Drug List:

Treprostini

Policy/Guideline:

Prescriber Specialty

This medication must be prescribed by or in consultation with a pulmonologist or cardiologist.

Coverage Criteria

Pulmonary Arterial Hypertension (PAH)¹⁻⁶

Authorization of 12 months may be granted for treatment of PAH when ALL of the following criteria are met:



AETNA BETTER HEALTH®
Coverage Policy/Guideline

Name: Treprostinil Page: 2 of 4

Effective Date: 10/15/2025 Last Review Date: 9/2025

| | | | |
|-------------|---|--|---|
| Applies to: | <input checked="" type="checkbox"/> Illinois | <input type="checkbox"/> Florida | <input checked="" type="checkbox"/> Florida Kids |
| | <input type="checkbox"/> New Jersey | <input checked="" type="checkbox"/> Maryland | <input type="checkbox"/> Michigan |
| | <input checked="" type="checkbox"/> Pennsylvania Kids | <input type="checkbox"/> Virginia | <input checked="" type="checkbox"/> Kentucky PRMD |

- Member has PAH defined as WHO Group 1 class of pulmonary hypertension (refer to Appendix).
- PAH was confirmed by either of the following:
 - Pretreatment right heart catheterization with all of the following results:
 - Mean pulmonary arterial pressure (mPAP) > 20 mmHg
 - Pulmonary capillary wedge pressure (PCWP) ≤ 15 mmHg
 - Pulmonary vascular resistance (PVR) > 2 Wood units. For pediatric members, pulmonary vascular resistance index (PVRI) > 3 Wood units x m² is also acceptable.
 - For infants less than one year of age, PAH was confirmed by Doppler echocardiogram if right heart catheterization cannot be performed.
- Patient is unable to take the required number of formulary alternatives (3) for the given diagnosis due to a trial and inadequate treatment response or intolerance, or a contraindication

Continuation of Therapy

Authorization of 12 months may be granted for members with an indication listed in the coverage criteria section who are currently receiving the requested medication through a paid pharmacy or medical benefit, and who are experiencing benefit from therapy as evidenced by disease stability or disease improvement.

Appendix

WHO Classification of Pulmonary Hypertension (PH)⁴

Note: Patients with heritable PAH or PAH associated with drugs and toxins might be long-term responders to calcium channel blockers.

Group 1: Pulmonary Arterial Hypertension (PAH)

- Idiopathic
 - Long-term responders to calcium channel blockers
- Heritable
- Associated with drugs and toxins
- Associated with:
 - Connective tissue disease
 - Human immunodeficiency virus (HIV) infection
 - Portal hypertension
 - Congenital heart disease
 - Schistosomiasis
- PAH with features of venous/capillary (pulmonary veno-occlusive disease [PVOD]/pulmonary capillary hemangiomatosis [PCH]) involvement



AETNA BETTER HEALTH®
Coverage Policy/Guideline

Name: Treprostinil

Page: 3 of 4

Effective Date: 10/15/2025

Last Review Date: 9/2025

| | | | |
|-------------|---|--|---|
| Applies to: | <input checked="" type="checkbox"/> Illinois | <input type="checkbox"/> Florida | <input checked="" type="checkbox"/> Florida Kids |
| | <input type="checkbox"/> New Jersey | <input checked="" type="checkbox"/> Maryland | <input type="checkbox"/> Michigan |
| | <input checked="" type="checkbox"/> Pennsylvania Kids | <input type="checkbox"/> Virginia | <input checked="" type="checkbox"/> Kentucky PRMD |

- Persistent PH of the newborn

Group 2: PH associated with Left Heart Disease

- Heart failure:
 - With preserved ejection fraction
 - With reduced or mildly reduced ejection fraction
 - Cardiomyopathies with specific etiologies (i.e., hypertrophic, amyloid, Fabry disease, and Chagas disease)
- Valvular heart disease:
 - Aortic valve disease
 - Mitral valve disease
 - Mixed valvular disease
- Congenital/acquired cardiovascular conditions leading to post-capillary PH

Group 3: PH associated with Lung Diseases and/or Hypoxia

- Chronic obstructive pulmonary disease (COPD) and/or emphysema
- Interstitial lung disease
- Combined pulmonary fibrosis and emphysema
- Other parenchymal lung diseases (i.e., parenchymal lung diseases not included in Group 5)
- Nonparenchymal restrictive diseases:
 - Hypoventilation syndromes
 - Pneumonectomy
- Hypoxia without lung disease (e.g., high altitude)
- Developmental lung diseases

Group 4: PH associated with Pulmonary Artery Obstructions

- Chronic thromboembolic PH
- Other pulmonary artery obstructions:
 - Sarcomas (high- or intermediate-grade or angiosarcoma)
 - Other malignant tumors (e.g., renal carcinoma, uterine carcinoma, germ-cell tumors of the testis)
 - Non-malignant tumors (e.g., uterine leiomyoma)
 - Arteritis without connective tissue disease
 - Congenital pulmonary artery stenoses
 - Hydatidosis



AETNA BETTER HEALTH®
Coverage Policy/Guideline

Name: Treprostinil Page: 4 of 4

Effective Date: 10/15/2025 Last Review Date: 9/2025

| | | | |
|-------------|---|--|---|
| Applies to: | <input checked="" type="checkbox"/> Illinois | <input type="checkbox"/> Florida | <input checked="" type="checkbox"/> Florida Kids |
| | <input type="checkbox"/> New Jersey | <input checked="" type="checkbox"/> Maryland | <input type="checkbox"/> Michigan |
| | <input checked="" type="checkbox"/> Pennsylvania Kids | <input type="checkbox"/> Virginia | <input checked="" type="checkbox"/> Kentucky PRMD |

Group 5: PH with Unclear and/or Multifactorial Mechanisms

- Hematological disorders, including inherited and acquired chronic hemolytic anemia and chronic myeloproliferative disorders
- Systemic disorders: Sarcoidosis, pulmonary Langerhans cell histiocytosis, and neurofibromatosis type 1
- Metabolic disorders, including glycogen storage diseases and Gaucher disease
- Chronic renal failure with or without hemodialysis
- Pulmonary tumor thrombotic microangiopathy
- Fibrosing mediastinitis
- Complex congenital heart disease

Approval Duration and Quantity Restrictions:

Approval: 12 months

References:

1. Treprostinil [package insert]. Princeton, NJ: Sandoz Inc.; April 2023.
2. Simonneau G, Montani D, Celermajer DS, et al. Haemodynamic definitions and updated clinical classification of pulmonary hypertension. *Eur Respir J.* 2019;53(1):1801913. doi:10.1183/13993003.01913-2018
3. Kovacs G, Bartolome S, Denton CP, et al. Definition, classification and diagnosis of pulmonary hypertension. *Eur Respir J.* 2024;64(4):2401324. doi: 10.1183/13993003.01324-2024
4. Chin KM, Gaine SP, Gerges C, et al. Treatment algorithm for pulmonary arterial hypertension. *Eur Respir J.* 2024;64(4):2401325. doi: 10.1183/13993003.01325-2024
5. Ivy D, Rosenzweig EB, Abman SH, et al. Embracing the challenges of neonatal and paediatric pulmonary hypertension. *Eur Respir J.* 2024;64(4):2401345. doi: 10.1183/13993003.01345-2024