



AETNA BETTER HEALTH®
Coverage Policy/Guideline

Name: Xolair-Omlyclo Page: 1 of 9

Effective Date: 9/5/2025 Last Review Date: 8/2025

Applies to:	<input type="checkbox"/> Illinois	<input type="checkbox"/> Florida	<input type="checkbox"/> Florida Kids
	<input checked="" type="checkbox"/> New Jersey	<input type="checkbox"/> Maryland	<input type="checkbox"/> Michigan
	<input type="checkbox"/> Pennsylvania Kids	<input type="checkbox"/> Virginia	<input type="checkbox"/> Kentucky PRMD

Intent:

The intent of this policy/guideline is to provide information to the prescribing practitioner outlining the coverage criteria for Xolair and Omlyclo under the patient's prescription drug benefit.

Description:

The indications below including FDA-approved indications and compendial uses are considered a covered benefit provided that all the approval criteria are met and the member has no exclusions to the prescribed therapy.

FDA-approved Indications^{1,2}

Allergic asthma

Indicated for adult and pediatric patients 6 years of age and older with moderate to severe persistent asthma who have a positive skin test or in vitro reactivity to a perennial aeroallergen and whose symptoms are inadequately controlled with inhaled corticosteroids. (*Reference Biologics in Severe Asthma New Jersey State Protocol.*)

Limitations of use

Not indicated for the relief of acute bronchospasm or status asthmaticus, or for treatment of other allergic conditions.

Chronic rhinosinusitis with nasal polyps (CRSwNP)

Indicated for add-on maintenance treatment of chronic rhinosinusitis with nasal polyps (CRSwNP) in adult patients 18 years of age and older with inadequate response to nasal corticosteroids.

IgE-mediated food allergy

Indicated for the reduction of allergic reactions (Type 1), including anaphylaxis, that may occur with accidental exposure to one or more foods in adult and pediatric patients aged 1 year and older with IgE-mediated food allergy.

Is to be used in conjunction with food allergen avoidance.

Limitations of use

Not indicated for the emergency treatment of allergic reactions, including anaphylaxis.

Chronic spontaneous urticaria (CSU)

Indicated for the treatment of adults and adolescents 12 years of age and older with chronic spontaneous urticaria (CSU) who remain symptomatic despite H1 antihistamine treatment.

Limitations of use

Not indicated for treatment of other forms of urticaria.



AETNA BETTER HEALTH®
Coverage Policy/Guideline

Name: Xolair-Omlyclo Page: 2 of 9

Effective Date: 9/5/2025 Last Review Date: 8/2025

Applies to:	<input type="checkbox"/> Illinois	<input type="checkbox"/> Florida	<input type="checkbox"/> Florida Kids
	<input checked="" type="checkbox"/> New Jersey	<input type="checkbox"/> Maryland	<input type="checkbox"/> Michigan
	<input type="checkbox"/> Pennsylvania Kids	<input type="checkbox"/> Virginia	<input type="checkbox"/> Kentucky PRMD

Compendial Uses¹²

- Immune checkpoint inhibitor-related toxicities
- Systemic mastocytosis

All other indications are considered experimental/investigational and not medically necessary.

Applicable Drug List:

Xolair
Omlyclo

Policy/Guideline:

Documentation

Submission of the following information is necessary to initiate the prior authorization review:

CRSwNP

Initial requests

- Chart notes or medical record documentation showing nasal endoscopy, anterior rhinoscopy, or computed tomography (CT) details (e.g., location, size), Meltzer Clinical Score, or endoscopic nasal polyp score (NPS) (where applicable).
- Chart notes, medical record documentation, or claims history supporting previous medications tried including drug, dose, frequency and duration. If therapy is not advisable, documentation of clinical reason to avoid therapy.

Continuation requests

Chart notes or medical record documentation supporting positive response to therapy.

IgE-Mediated Food Allergy

Initial requests

Chart notes, medical record documentation, or laboratory tests showing the following (where applicable):

- Pre-treatment allergen-specific IgE level.
- Skin-prick test wheal diameter.
- Pre-treatment serum IgE level.
- Positive result of a physician controlled oral food challenge.
- History of a systemic reaction to a food.



AETNA BETTER HEALTH®
Coverage Policy/Guideline

Name: Xolair-Omlyclo Page: 3 of 9

Effective Date: 9/5/2025 Last Review Date: 8/2025

Applies to:	<input type="checkbox"/> Illinois	<input type="checkbox"/> Florida	<input type="checkbox"/> Florida Kids
	<input checked="" type="checkbox"/> New Jersey	<input type="checkbox"/> Maryland	<input type="checkbox"/> Michigan
	<input type="checkbox"/> Pennsylvania Kids	<input type="checkbox"/> Virginia	<input type="checkbox"/> Kentucky PRMD

Continuation requests

Chart notes or medical record documentation supporting positive response to therapy (e.g., reduction or absence of hypersensitivity to food allergen).

CSU

Initial requests

Chart notes, medical record documentation, or claims history supporting previous medications tried, including response to therapy.

Continuation requests

Chart notes or medical record documentation supporting positive response to therapy.

Immune Checkpoint Inhibitor-Related Toxicity

Initial requests

Chart notes or medical record documentation showing pre-treatment IgE level.

Continuation requests

Chart notes or medical record documentation supporting positive response to therapy.

Systemic Mastocytosis

Initial requests

- Chart notes or medical record documentation supporting diagnosis of systemic mastocytosis.
- Chart notes, medical record documentation, or claims history supporting previous medications tried (where applicable).

Continuation requests

Chart notes or medical record documentation supporting positive response to therapy.

Prescriber Specialties

This medication must be prescribed by or in consultation with one of the following:

- CRSwNP: allergist/immunologist or otolaryngologist
- IgE-mediated food allergy: allergist/immunologist
- Chronic spontaneous urticaria: allergist/immunologist or dermatologist
- Immune checkpoint inhibitor-related toxicity: dermatologist, hematologist, or oncologist



AETNA BETTER HEALTH®
Coverage Policy/Guideline

Name: Xolair-Omylclo Page: 4 of 9

Effective Date: 9/5/2025 Last Review Date: 8/2025

Applies to:	<input type="checkbox"/> Illinois	<input type="checkbox"/> Florida	<input type="checkbox"/> Florida Kids
	<input checked="" type="checkbox"/> New Jersey	<input type="checkbox"/> Maryland	<input type="checkbox"/> Michigan
	<input type="checkbox"/> Pennsylvania Kids	<input type="checkbox"/> Virginia	<input type="checkbox"/> Kentucky PRMD

Coverage Criteria

Chronic Rhinosinusitis with Nasal Polyps (CRSwNP)^{1,2,10,11,15-17}

Authorization of 6 months may be granted for adult members who have previously received a biologic drug indicated for chronic rhinosinusitis with nasal polyps (CRSwNP) in the past year.

Authorization of 6 months may be granted for treatment of CRSwNP when all of the following criteria are met:

- Member is 18 years of age or older.
- Member has bilateral nasal polyps and chronic symptoms of sinusitis despite intranasal corticosteroid treatment for at least 4 weeks unless contraindicated or not tolerated.
- Member has one of the following:
 - A bilateral nasal endoscopy, anterior rhinoscopy, or computed tomography (CT) showing polyps reaching below the lower border of the middle turbinate or beyond in each nostril.
 - Meltzer Clinical Score of 2 or higher in both nostrils.
 - A total endoscopic nasal polyp score (NPS) of at least 5 with a minimum score of 2 for each nostril.
- Member has symptoms of nasal blockage, congestion, or obstruction plus one of the following additional symptoms:
 - Rhinorrhea (anterior/posterior).
 - Reduction or loss of smell.
 - Facial pain or pressure.
- Member will continue to use a daily intranasal corticosteroid while being treated with the requested medication, unless contraindicated or not tolerated.

IgE-Mediated Food Allergy^{1,2,19,20}

Authorization of 6 months may be granted for reduction of IgE-mediated food allergy reactions when all of the following criteria are met:

- Member is 1 year of age or older.
- IgE-mediated food allergy has been confirmed by either of the following:
 - Pre-treatment food allergen-specific IgE level greater than or equal to 6 IU/mL.
 - Skin-prick test (SPC) with wheal diameter greater than or equal to 4 mm.
- Member has either of the following:
 - A positive physician controlled oral food challenge (e.g., moderate to severe skin, respiratory, or gastrointestinal [GI] symptoms).



AETNA BETTER HEALTH®
Coverage Policy/Guideline

Name: Xolair-Omlyclo Page: 5 of 9

Effective Date: 9/5/2025 Last Review Date: 8/2025

Applies to:	<input type="checkbox"/> Illinois	<input type="checkbox"/> Florida	<input type="checkbox"/> Florida Kids
	<input checked="" type="checkbox"/> New Jersey	<input type="checkbox"/> Maryland	<input type="checkbox"/> Michigan
	<input type="checkbox"/> Pennsylvania Kids	<input type="checkbox"/> Virginia	<input type="checkbox"/> Kentucky PRMD

- History of a systemic reaction to a food.
- Member has a pre-treatment serum IgE level greater than or equal to 30 IU/mL.
- Member will continue to follow a food-allergen avoidance diet.

Chronic Spontaneous Urticaria^{1,2,7,8}

Authorization of 6 months may be granted for members 12 years of age or older who have previously received a biologic drug indicated for chronic spontaneous urticaria in the past year.

Authorization of 6 months may be granted for treatment of chronic spontaneous urticaria when all of the following criteria are met:

- Member is 12 years of age or older.
- Member remains symptomatic despite treatment with up-dosing (in accordance with EAACI/GA2LEN/EuroGuiDerm/APAAACI guidelines) of a second-generation H1 antihistamine (e.g., cetirizine, fexofenadine, levocetirizine, loratadine) for at least 2 weeks.
- Member has been evaluated for other causes of wheals (hives) and/or angioedema, including bradykinin-related angioedema and interleukin-1-associated urticarial syndromes (auto-inflammatory disorders, urticarial vasculitis).
- Member has experienced a spontaneous onset of wheals (hives), angioedema, or both, for at least 6 weeks.

Immune Checkpoint Inhibitor-Related Toxicity^{12,21}

Authorization of 12 months may be granted for treatment of immune checkpoint inhibitor-related toxicity when both of the following criteria are met:

- The member has a refractory case of immune-therapy related severe (G3) pruritus with no response to gabapentinoids in one month.
- The member has elevated IgE levels.

Systemic Mastocytosis^{12,13}

Authorization of 12 months may be granted for treatment of systemic mastocytosis when both of the following criteria are met:

- The major and at least one minor diagnostic criterion for systemic mastocytosis are present or three or more minor diagnostic criteria are present (see Appendix).
- The requested medication will be used in any of the following treatment settings:



AETNA BETTER HEALTH®
Coverage Policy/Guideline

Name: Xolair-Omlyclo Page: 6 of 9

Effective Date: 9/5/2025 Last Review Date: 8/2025

Applies to:	<input type="checkbox"/> Illinois	<input type="checkbox"/> Florida	<input type="checkbox"/> Florida Kids
	<input checked="" type="checkbox"/> New Jersey	<input type="checkbox"/> Maryland	<input type="checkbox"/> Michigan
	<input type="checkbox"/> Pennsylvania Kids	<input type="checkbox"/> Virginia	<input type="checkbox"/> Kentucky PRMD

- Used as stepwise prophylactic treatment for chronic mast cell mediator-related cardiovascular and pulmonary symptoms when the member has tried both of the following:
 - H1 blockers and H2 blockers
 - Corticosteroids
- Used for prevention of recurrent unprovoked anaphylaxis.
- Used for prevention of hymenoptera or food-induced anaphylaxis, with negative specific IgE or negative skin test.
- Used to improve tolerability of venom immunotherapy.

Continuation of Therapy

Chronic Rhinosinusitis with Nasal Polyps (CRSwNP)^{1,2,10,11,15}

Authorization of 12 months may be granted for continuation of treatment of CRSwNP when all of the following criteria are met:

- Member is 18 years of age or older.
- Member has experienced a positive response as evidenced by improvement in signs and symptoms (e.g., improvement in nasal congestion, nasal polyp size, loss of smell, anterior or posterior rhinorrhea, sino-nasal inflammation, hyposmia and/or facial pressure or pain or reduction in corticosteroid use).
- Member will continue to use a daily intranasal corticosteroid while being treated with the requested medication, unless contraindicated or not tolerated.

IgE-Mediated Food Allergy^{1,2,20}

Authorization of 12 months may be granted for reduction of IgE-mediated food allergy reactions when all of the following criteria are met:

- Member is 1 year of age or older.
- Member has achieved or maintained a positive clinical response to therapy as evidenced by a reduction or absence of hypersensitivity (e.g., moderate to severe skin, respiratory or GI symptoms) to food-allergen.
- Member will continue to maintain a food-allergen avoidance diet.

Chronic Spontaneous Urticaria^{1,2,7}

Authorization of 12 months may be granted for continuation of treatment of chronic spontaneous urticaria when all of the following criteria are met:

- Member is 12 years of age or older.



AETNA BETTER HEALTH®
Coverage Policy/Guideline

Name:	Xolair-Omlyclo	Page:	7 of 9
Effective Date:	9/5/2025	Last Review Date:	8/2025
Applies to:	<input type="checkbox"/> Illinois <input checked="" type="checkbox"/> New Jersey <input type="checkbox"/> Pennsylvania Kids	<input type="checkbox"/> Florida <input type="checkbox"/> Maryland <input type="checkbox"/> Virginia	<input type="checkbox"/> Florida Kids <input type="checkbox"/> Michigan <input type="checkbox"/> Kentucky PRMD

- Member has experienced a positive response (e.g., improved symptoms, decrease in weekly urticaria activity score [UAS7]) since initiation of therapy.

Immune Checkpoint Inhibitor-Related Toxicity and Systemic Mastocytosis^{12,21}

Authorization of 12 months may be granted for continuation of treatment of refractory immune checkpoint inhibitor-related severe (G3) pruritus or systemic mastocytosis when the member achieves or maintains a positive clinical response as evidenced by low disease activity or improvement in signs and symptoms of the condition.

Other

For all indications: Member cannot use the requested medication concomitantly with any other biologic drug or targeted synthetic drug for the same indication.

Note: If the member is a current smoker or vaper, they should be counseled on the harmful effects of smoking and vaping on pulmonary conditions and available smoking and vaping cessation options.

Appendix

2024 WHO Diagnostic Criteria for Systemic Mastocytosis¹³

- Major Criteria: multifocal, dense infiltrates of mast cells (at least 15 mast cells in aggregates) detected in sections of bone marrow and/or other extracutaneous organs
- Minor Criteria
 - Atypical mast cell morphology, including spindle shape or immature morphology, present in greater than 25% of all mast cells on bone marrow smears or in other extracutaneous organ(s)
 - KIT p.D816V mutation or other activating KIT mutation(s) detected in peripheral blood, bone marrow, or other extracutaneous organ(s)
 - Mast cells aberrantly express one or more of the following antigens: CD2, CD25, CD30
 - Baseline serum tryptase concentration of greater than 20 ng/mL in the absence of an associated myeloid neoplasm; in the case of a known HαT, the tryptase level could be adjusted

Approval Duration and Quantity Restrictions:

Approval:

- Initial approval:
 - Chronic Spontaneous Urticaria: 6 – 12 months



AETNA BETTER HEALTH®
Coverage Policy/Guideline

Name:	Xolair-Omlyclo	Page:	8 of 9
Effective Date:	9/5/2025	Last Review Date:	8/2025
Applies to:	<input type="checkbox"/> Illinois <input checked="" type="checkbox"/> New Jersey <input type="checkbox"/> Pennsylvania Kids	<input type="checkbox"/> Florida <input type="checkbox"/> Maryland <input type="checkbox"/> Virginia	<input type="checkbox"/> Florida Kids <input type="checkbox"/> Michigan <input type="checkbox"/> Kentucky PRMD

- Nasal Polyps: 6 – 12 months
- Immune Checkpoint Inhibitor-related toxicities: 12 months
- Systemic Mastocytosis: 12 months
- Continuation: 12 months

Quantity Level Limit:

- Xolair 150 mg vial: 8 vials per 28 days
- Xolair 75 mg single-dose prefilled syringe: 2 syringes per 28 days
- Xolair 150 mg single-dose prefilled syringe: 8 syringes per 28 days
- Xolair (omalizumab) 300 mg/2 mL single-dose prefilled syringe/autoinjector: 4 syringes/autoinjectors per 28 days
- Omlyclo (omalizumab-igec) 75 mg/0.5 mL single-dose prefilled syringe: 2 syringes per 28 days
- Omlyclo (omalizumab-igec) 150 mg/1 mL single-dose prefilled syringe: 8 syringes per 28 days

References:

1. Xolair [package insert]. South San Francisco, CA: Genentech, Inc.; February 2024.
2. Omlyclo [package insert]. Jersey City, NJ: Celltrion USA, Inc.; March 2025.
3. National Institutes of Health. National Asthma Education and Prevention Program Expert Panel Report 3: Asthma Management Guidelines: Focused Updates 2020. Bethesda, MD: National Heart Lung and Blood Institute; December 2020. Available at <https://www.nhlbi.nih.gov/sites/default/files/publications/AsthmaManagementGuidelinesReport-2-4-21.pdf>. Accessed March 1, 2025.
4. Global Initiative for Asthma (GINA). Global Strategy for Asthma Management and Prevention. 2024 update. Available at: https://ginasthma.org/wp-content/uploads/2024/05/GINA-2024-Strategy-Report-24_05_22_WMS.pdf. Accessed March 1, 2025.
5. Strunk RC, Bloomberg GR. Omalizumab for asthma. N Engl J Med. 2006;354(25):2689-2695.
6. Kew KM, Karner C, Mindus SM. Combination formoterol and budesonide as maintenance and reliever therapy versus combination inhaler maintenance for chronic asthma in adults and children (review). Cochrane Database Syst Rev. 2013;12:CD009019.
7. Zuberbier T, Abdul Latiff AH, Abuzakouk M, et al. The international EAACI/GA²LEN/EuroGuiDerm/APAAACI guideline for the definition, classification, diagnosis, and management of urticaria. Allergy. 2022 Mar;77(3):734-766.
8. Bernstein DI, Blessing-Moore J, Cox L, et al. The diagnosis and management of acute and chronic urticaria: 2014 update. American Academy of Allergy, Asthma & Immunology Practice Parameter. <http://www.aaaai.org/practice-resources/statements-and-practice-parameters/practice-parameter-guidelines.aspx>. Accessed March 1, 2025.



AETNA BETTER HEALTH®
Coverage Policy/Guideline

Name:	Xolair-Omlyclo	Page:	9 of 9
Effective Date:	9/5/2025	Last Review Date:	8/2025
Applies to:	<input type="checkbox"/> Illinois <input checked="" type="checkbox"/> New Jersey <input type="checkbox"/> Pennsylvania Kids	<input type="checkbox"/> Florida <input type="checkbox"/> Maryland <input type="checkbox"/> Virginia	<input type="checkbox"/> Florida Kids <input type="checkbox"/> Michigan <input type="checkbox"/> Kentucky PRMD

9. Maurer M, Rosen K, Hsieh HJ, et al. Omalizumab for the treatment of chronic idiopathic or spontaneous urticaria. *N Engl J Med.* 2013;368(10):924-935.
10. ClinicalTrials.gov. National Library of Medicine (US). Identifier NCT03280550, A Clinical Trial of Omalizumab in Participants with Chronic Rhinosinusitis with Nasal Polyps (POLYP 1) 2017 Sep 12. Available from: <https://clinicaltrials.gov/ct2/show/NCT03280550>.
11. ClinicalTrials.gov. National Library of Medicine (US). Identifier NCT03280537, A Clinical Trial of Omalizumab in Participants with Chronic Rhinosinusitis with Nasal Polyps (POLYP 2) 2017 Sep 12. Available from: <https://clinicaltrials.gov/ct2/show/NCT03280537>.
12. The NCCN Drugs & Biologics Compendium® © 2025 National Comprehensive Cancer Network, Inc. Available at: <http://www.nccn.org>. Accessed March 1, 2025.
13. National Comprehensive Cancer Network. NCCN Clinical Practice Guidelines in Oncology: Systemic Mastocytosis. Version 1.2025. https://www.nccn.org/professionals/physician_gls/pdf/mastocytosis.pdf. Accessed March 1, 2025.
14. Cloutier MM, Dixon AE, Krishnan JA, et al. Managing asthma in adolescents and adults: 2020 asthma guideline update from the National Asthma Education and Prevention Program. *JAMA.* 2020;324(22): 2301-2317.
15. Bachert C, Han JK, Wagenmann M, et al. EUFOREA expert board meeting on uncontrolled severe chronic rhinosinusitis with nasal polyps (CRSwNP) and biologics: Definitions and management. *J Allergy Clin Immunol.* 2021;147(1):29-36.
16. Fokkens WJ, Lund VJ, Hopkins C, et al. European Position Paper on Rhinosinusitis and Nasal Polyps 2020. *Rhinology.* 2020;58(Suppl S29):1-464.
17. Hopkins C. Chronic Rhinosinusitis with Nasal Polyps. *N Engl J Med.* 2019;381(1):55-63.
18. Rank MA, Chu DK, Bognanni A, et al. The Joint Task Force on Practice Parameters GRADE guidelines for the medical management of chronic rhinosinusitis with nasal polyposis. *J Allergy Clin Immunol.* 2023 Feb;151(2):386-398.
19. NIAID-Sponsored Expert Panel, Boyce JA, Assa'ad A, et al. Guidelines for the diagnosis and management of food allergy in the United States: report of the NIAID-sponsored expert panel. *J Allergy Clin Immunol.* 2010;126(6 Suppl):S1-58.
20. Wood RA, Togias A, Sicherer SH, et al. Omalizumab for the Treatment of Multiple Food Allergies. *N Engl J Med.* 2024;390(10):889-899.
21. NCCN Clinical Practice Guidelines in Oncology® (NCCN Guidelines®). Management of Immune Checkpoint-Related Toxicities. Version 1.2025. Available at: www.nccn.org. Accessed March 1, 2025.