Pharmacy Prior Authorization

AETNA BETTER HEALTH ILLINOIS FAMILY HEALTH PLAN (MEDICAID)

Epogen-Procrit (Medicaid)

This fax machine is located in a secure location as required by HIPAA regulations.

Complete/review information, sign and date. Fax signed forms to Aetna Better Health Illinois at 1-844-242-0908.

When conditions are met, we will authorize the coverage of Epogen-Procrit (Medicaid).

Please note that all authorization requests will be reviewed as the AB rated generic (when available) unless states otherwise.

| Drug Name (please circle) | | | |
|---|---|---|---|
| Epogen (epoetin alfa) | Procrit (epoetin alfa) | | |
| Other, please specify | | | |
| Quantity | Frequency Strength _ | | |
| Route of Administration | Expected Length of therapy | | |
| Patient Information | | | |
| Patient Name: | | | |
| Patient ID: | | | |
| | | | |
| | | | |
| Patient Phone: | | | |
| Prescribing Physician | | | |
| Physician Name: | | | |
| Specialty: | NPI Number: | | |
| Physician Fax: | Physician Phone: | | |
| Physician Address: | City, State, Zip: | | |
| Diagnosis: | ICD Code: | | |
| Please circle the appropriate answer | for each question. | | |
| Has this plan authorized the previous authorization is contact. | his medication in the past for this patient (i.e., on file under this plan)? | Υ | N |
| [If no, skip to question 3.] | | | |
| Hemoglobin less than 11 ghas adequate iron stores t | h of the following conditions for approval: A) g/dL within the last 2 weeks, and B) Patient to support erythropoiesis (e.g., serum ferritin rin saturation above 20%) | Υ | N |
| Please document hemoglodrawn: | obin and results of iron studies including date | | |

Reference Number: C4916-A / Effective Date: 06/16/2017

| | [No further questions.] | | |
|----|---|---|---|
| 3. | Does the patient have adequate iron stores to support erythropoiesis as evidenced by one of the following: A) Serum ferritin greater than or equal to100 ng/ml and transferrin saturation (iron saturation) greater than or equal to 20%, or B) Normal serum iron, TIBC and serum ferritin, or C) Reticulocyte hemoglobin content (CHr) greater than 29 | Y | N |
| | Please document Iron Studies obtained, results, and date drawn: | | |
| | [If no, then no further questions.] | | |
| 4. | Does the patient have uncontrolled high blood pressure? | Υ | N |
| | [If yes, then no further questions.] | | |
| 5. | Does the patient have a diagnosis of anemia due to chronic kidney disease? | Υ | N |
| | [If no, skip to question 7.] | | |
| 6. | Does the patient have hemoglobin less than 10 g/dL within 2 weeks prior to initiating therapy? | Υ | N |
| | Please document hemoglobin and date drawn: | | |
| | [If no, then no further questions.] | | |
| | [If yes, skip to question 18.] | | |
| 7. | Is therapy requested for the treatment of anemia in a cancer patient? | Υ | N |
| | [If no, skip to question 10.] | | |
| 8. | Is the patient currently receiving chemotherapy? | Υ | N |
| | [If no, then no further questions.] | | |
| 9. | Does the patient meet all of the following conditions for approval: A) Hemoglobin less than 10 g/dL within the 2 weeks prior to starting therapy, B) Diagnosis of non-myeloid malignancy (e.g., solid tumor), and C) Patient will receive chemotherapy for at least 2 additional months | Υ | N |

Reference Number: C4916-A / Effective Date: 06/16/2017

10. [If yes, go to question 18.]

[If no, then no further question]

Please document hemoglobin and date drawn:

Ν

| 11. Is the request for a patient with high risk factors for bleeding who will be undergoing elective, noncardiac, and nonvascular surgery? | Y | N |
|--|---|---|
| [If no, skip to question 14.] | | |
| 12. Does the patient have a hemoglobin level greater than 10 but less than or equal to 13 g/dL within 30 days prior to the planned surgery? | Υ | N |
| Please document hemoglobin and date drawn: | | |
| [If no, then no further questions.] | | |
| 13. Is this request for Procrit? | Υ | N |
| [If no, then no further questions.] | | |
| 14. Has the patient experienced treatment failure or intolerable side effects with Epogen? | Υ | N |
| [No further questions.] | | |
| 15. Is therapy requested for the treatment of anemia in a patient with HIV who is taking zidovudine? | Υ | N |
| [If no, skip to question 16.] | | |
| 16. Is the zidovudine dose less than or equal to 4200 mg/week? | Υ | N |
| (Note: it is recommended to decrease the dose of zidovudine to 4200 mg per week or less if the patient is experiencing anemia.) | | |
| [If no, then no further questions.] | | |
| [If yes, skip to question 17.] | | |
| 17. Is therapy requested for the treatment of anemia associated with myelodysplastic syndrome (MDS)? | Υ | N |
| [If no, then no further questions.] | | |
| 18. Does the patient meet all of the following conditions for approval: A) Hemoglobin is less than 10 g/dL within 2 weeks prior to initiating therapy, and B) Erythropoietin level is less than or equal to 500 IU/L | Υ | N |
| Please document erythropoietin and hemoglobin levels and dates drawn: | Υ | N |
| [If no, then no further questions.] | | |

Reference Number: C4916-A / Effective Date: 06/16/2017

| Proscribor (Or Authorized) Signature | 0 | |
|--|---|---|
| affirm that the information given on this form is true and accurate as of this date. | | |
| Comments: | | |
| 20. Has the patient experienced treatment failure or intolerable side effects with Epogen? | Υ | N |
| [If no, then no further questions.] | | |
| 19. Is this request for Procrit? | Υ | N |

Reference Number: C4916-A / Effective Date: 06/16/2017